


# The Norwegian Zoonoses Report 2019



# The Norwegian Zoonoses Report 2019

## Content

Summary .....	2
Introduction .....	2
Origin of data .....	2
Preventive and protective measures .....	3
Acknowledgements .....	4
Salmonellosis .....	5
Campylobacteriosis .....	7
Yersiniosis .....	9
Listeriosis .....	10
Verotoxin producing <i>E. coli</i> (VTEC) .....	11
Tuberculosis .....	12
Brucellosis .....	13
Trichinellosis .....	14
Echinococcosis .....	15
Toxoplasmosis .....	16
Rabies .....	17
Q-fever .....	18
BSE and vCJD .....	19
Antimicrobial resistance .....	20
Foodborne outbreaks .....	21
Appendix Tables 2018 .....	22

### Authors

Hannah J. Jørgensen<sup>1</sup>, Kjell Hauge<sup>2</sup>, Heidi Lange<sup>3</sup>, Trude Marie Lyngstad<sup>3</sup>, Berit Tafjord Heier<sup>1</sup>

<sup>1</sup>Norwegian Veterinary Institute

<sup>2</sup>Norwegian Food Safety Authority

<sup>3</sup>Norwegian Institute of Public Health

In collaboration with



ISSN 1890-3290

© Norwegian Veterinary Institute 2020

Design Cover: Reine Linjer  
Photo front page: Colourbox

## Summary

In general, the occurrence of most zoonoses in 2019 in Norway was similar compared to previous years.

In humans the incidence of zoonotic disease has remained stable; the exception was *E. coli* (VTEC) for which the number of cases has continued to rise. Although the increase observed in recent years can be explained by changes in diagnostic methods, the trend is still of concern because the infection can cause serious disease.

## Introduction

The Zoonosis Report is published annually in Norway in accordance with the requirements of the EU Council Directive 2003/99/EC. In addition, data on specified zoonoses in feed, animals and food are reported to the European Food Safety Authority (EFSA). Corresponding data from humans are reported to the European Center for Disease Control (ECDC). These two European institutions compile an annual European zoonosis report based on the received data:

([http://www.efsa.europa.eu/en/publications/advanced-search/?sub\\_subject=61616](http://www.efsa.europa.eu/en/publications/advanced-search/?sub_subject=61616)).

The Norwegian Veterinary Institute (NVI) is responsible for reporting of Norwegian data to EFSA, while the Norwegian Institute of Public Health (NIPH) reports Norwegian data to ECDC. The Zoonosis Report is written by the NVI in collaboration with the Norwegian Food Safety Authority (NFSA) and NIPH.

## Origin of data

### *Humans*

“The Norwegian Surveillance System for Communicable Diseases” (MSIS) was implemented nationally in Norway in 1975, and the NIPH is responsible for managing the system. The main purpose of MSIS is surveillance to describe trends and detect outbreaks of communicable diseases.

According to the Infectious Disease Control Act, all clinicians and laboratories that analyse samples from humans must report all cases of specified communicable diseases (at present 65 different diseases). All zoonoses described in this report, with the exception of toxoplasmosis, are notifiable.

Patients who have not travelled abroad during the incubation period for the diagnosed infection are classified as “infected in Norway”. Patients who develop the diagnosed infection abroad or shortly after returning home to Norway are classified as “infected abroad”. Patients for whom information regarding travel is not available are classified as «unknown origin» with respect to where the infection was contracted.

The District Medical Officer must notify the NFSA in cases where humans are believed to be infected from animals or food.

### *Feed, animals and food*

The data presented in this report are obtained through national surveillance programmes, projects, diagnostic investigations and various inspections performed by public authorities and private companies. Two types of data are reported:

- Data on detected notifiable diseases (reported to the NFSA) and data from public surveillance. The NFSA decides which infectious agents are notifiable and which surveillance programmes should be carried out. The NVI assists with planning and laboratory analyses, data processing and reporting. Testing of animals and food for various zoonotic agents is also in association with import and export. In addition, surveillance in association with commercial slaughter through pre-and post-mortem inspections are carried out by the NFSA.
- Data from diagnostic investigations and data from internal control systems of food-, and feed-producing companies are also included in the Zoonosis Report. All laboratories are obliged to report any detection of notifiable diseases in animals to the NFSA. A large proportion of the laboratory diagnostics (including pathology) performed on animals in Norway is performed by the NVI. In cases where laboratories abroad are used, the responsible veterinarian is obliged to report any detection of notifiable disease in animals. Data from internal control of companies are not always available. One exception is Salmonella-control in feed producing companies, where data from most of the performed internal control is made available and is presented in this report.

Notifiable diseases/agents in animal and humans are presented in Table 1.

### **Preventive and protective measures**

Norway has strict regulations to prevent introduction and spread of certain infections in animals and humans.

#### *Humans*

When clusters of notifiable zoonoses are detected in humans, investigations are performed to trace the source of infection and measures to prevent new cases are implemented. In cases where food or animals are suspected to be the source, the NFSA is notified.

People employed in the food industry should not work while symptomatic with infections that may be transmitted through food. Before returning to work they should have two negative faecal samples after clinical improvement. For EHEC/VTEC/*S. Typhi*/*S. Paratyphi*/*S. dysenteriae* 1 the number of negative faecal samples should be three.

#### *Feed, animals and food*

According to the Food Act (Matloven), Food Business Operators are responsible for implementing appropriate measures to prevent the occurrence or spread of contagious disease in animals, and to notify the NFSA about any suspicion of contagious disease in animals that has potential to cause significant negative consequences for society.

The Regulation on Notification of Diseases in Animals states that veterinarians and laboratories must notify the NFSA about specified animal diseases categorized as A-, B-, and C-diseases. In addition, there is a general duty to notify diseases in animals that:

- could cause death or serious disease in humans.
- could result in high numbers of animals becoming diseased or exposed to infection.
- could result in substantial economic losses for society.
- could cause other substantial consequences for society.
- are presumed not to exist in Norway or have an unexpected distribution.
- compromise animal health in an unexpected manner or in an unexpected fashion.

Suspicion or diagnosis of Group A and B diseases in animals should be notified immediately to the Norwegian Food Safety Authority. Diagnosis of group C disease in animals should be reported to the Food Safety Authority within seven working days.

If a group A- or B-disease is detected in animals in Norway, restrictions will be imposed on the infected animal or animal holding, and efforts will be made to eradicate the infective agent. The imposed/recommended measures depend on animal species, management system, and the infective agent. In cases where a zoonosis is detected or suspected, the NFSA must notify the District Medical Officer if the infection has transmitted - or may transmit - to humans.

Companies that produce or sell food are themselves responsible for ensuring that the products they produce or sell are safe to consume. The NFSA follows up and inspects the food industry facilities to ensure that they exercise their responsibility. Food producers must also consider zoonoses in their internal control systems. In addition to the national surveillance programmes and various short-term projects initiated by the head office of the NFSA, the regional offices of the NFSA perform some sampling. However, the data from regional offices are not included in this report.

In total, 14 border inspection posts and 7 associated control centres in Norway perform control of foods and foodstuffs of animal origin that are imported from non EU and non-EEA-countries.

If a zoonotic agent is detected in a food or foodstuff, measures are carried out to prevent spread and to identify the source. The District Medical Officer must be notified, and if there is a risk that animals have been infected or may become infected, the NFSA must perform further investigations.

Table 1. Disease/agents included in the zoonosis report in 2019 and their status with respect to notifiability and existing surveillance programmes.

Disease/agent	Notifiability			Feed, animals and food
	Humans	Feed and food	Animals	Surveillance programme
Salmonellosis	Yes	Yes	Yes (B-disease)	Yes
Campylobacteriosis	Yes	No*	No**	Yes
Yersiniosis	Yes	No*	No	No (occasionally)
Listeriosis	Yes	No*	Yes (C-disease)	No
Pathogenic <i>E. coli</i>	Yes	Yes*	Yes*	No (occasionally)
Tuberculosis	Yes	Yes	Yes (B-disease)	Yes
Brucellosis	Yes	Yes	Yes (A-disease)	Yes
Trichinellosis	Yes	Yes	Yes (B-disease)	Yes
Echinococcosis	Yes	Yes	Yes (B-disease)	Yes
Toxoplasmosis	No	No	Yes (C-disease)	No
Rabies	Yes	-	Yes (A-disease)	No
Q-fever	Yes	-	Yes (C-disease)	No (occasionally)
BSE og vCJD	Yes	-	Yes (B-disease)	Yes

\* Some conditions are notifiable according to national regulation within specific areas. Otherwise, the food law contains a general obligation to immediately inform the competent authorities if there exists a risk or potential risk (to human, animal and plant health) of significant consequences to the society.

\*\* The exception is broiler chickens slaughtered before 51 days of age between May and October, because these are included in the surveillance programme, and measures are implemented if samples are positive.

## Acknowledgements

Institute of Marine Research, Geno, Norsvin and the feed industry are acknowledged for providing data for this report.

## Salmonellosis

### The disease and its transmission routes

There are more than two thousand variants of *Salmonella* bacteria. The most common symptom of infection is diarrhoea, both in humans and in animals, but healthy carriage is not uncommon. *Salmonella* are shed in faeces and the most important sources of infection are contaminated food, feed or water. *Salmonella* can also spread through direct contact with infected individuals.

### Surveillance and control

Salmonellosis in humans is notifiable in Norway. From 2017, both *Salmonella* infections verified by PCR and/or by culture are registered in MSIS. *Salmonella*-infection in animals is notifiable (group B-disease). Detection of *Salmonella* in feed or food must also be reported to the NFSA.

Surveillance of *Salmonella* in feed, cattle, swine and poultry (live animals and animal products) started in 1995. Testing is performed in cases of disease, live animal import and as part of Salmonella control systems in feed production. Vaccination of animals against *Salmonella* is forbidden in Norway.

## Results 2019

The number of reported cases of salmonellosis in humans (1094) increased slightly in 2019 compared to the previous four years (Figure 1). Information on the most frequently detected serotypes is presented in the Appendix.




Figure 1. Reported cases of salmonellosis in humans. Data from MSIS.

There is a surveillance programme for *Salmonella* in live animals. One broiler flock and one turkey production flock were positive, giving an estimated *Salmonella* prevalence of 0.04% in poultry flocks for slaughter. A total of 3,419 lymph node samples from pigs were examined, and two samples from slaughter pigs were positive for *Salmonella* giving an estimated prevalence of 0.06% at the individual carcass level. From cattle, a total of 3,151 lymph node samples were examined, and *Salmonella* was not detected.

In 2019, only a few cases of salmonellosis were diagnosed in cats (n=2) dogs (n=1), and horses (n=2), but out of 19 samples from reptiles, 16 samples were positive for *Salmonella*. Details on *Salmonella* testing of feed, animals and food are shown in the Appendix.

## Evaluation of the current situation

The number of salmonellosis cases in humans has decreased over the past 10 years. The number of cases infected in Norway has remained relatively stable with an incidence of 3.4 to 4.8 in the last 5 years (2014-2018) but increased to 6.0 in 2019 mainly due to a national outbreak of *Salmonella* Agbeni which included 56 cases. 59% of the human cases reported to have contracted the infection abroad. The reduced prevalence of *Salmonella* in European poultry is presumed to contribute to the observed reduction. Data from outbreaks of salmonellosis indicate that a great variety of foods can be implicated. When infection is contracted in Norway, imported foods are more often implicated than foods produced in Norway.

In Norway, food-producing animals are very rarely infected with *Salmonella*. This is well documented in the surveillance program (Figure 2). *Salmonella diarizonae* is occasionally detected in Norwegian sheep (n=5 flocks in 2019). This variant is only rarely associated with disease in animals and is not considered a public health threat.




Figure 2. The number of positive samples in the *Salmonella* surveillance programme.

*Salmonella* is occasionally detected in dogs and cats and in reptiles in Norway, but the number of detected cases in dogs and cats has decreased slightly in the last few years. A reoccurrence of a monophasic *Salmonella* Typhimurium at an equine hospital resulted in two infected horses in 2019. The hospital had an outbreak with the same strain in 2018, and rigorous testing of hospitalized horses revealed the reoccurrence early. Nevertheless, a large number of contact horses (>200) were tested as part of the follow up. In the last years only a few cases of salmonellosis in animals are detected annually in the diagnostic services at the NVI, and only a few cases are reported to the Food Safety Authority.

In 2017, an exemption was made for 19 species on the general ban on import and marketing of reptiles in Norway. Reptiles frequently carry *Salmonella* and may pose a source of infection to humans. *Salmonella* Typhimurium can sometimes be detected from wild birds and hedgehogs in Norway. Contamination of food and water by these animals may lead to infection of humans.

Feed given to domestic animals in Norway is generally free from *Salmonella* in Norway, but *Salmonella* is sometimes detected in feed factories, especially those producing fish feed. Continued surveillance of *Salmonella* in animals, feed and food is necessary for early detection, to facilitate control and to sustain the beneficial situation with respect to *Salmonella* in Norway.

## Campylobacteriosis

### The disease and its transmission routes

There are many *Campylobacter* variants, but *C. jejuni* and *C. coli* are the most important zoonoses. These are commonly found in the guts of healthy birds, and humans may contract the infection through contaminated food or water or by direct contact. Diarrhoea is the most common symptom of campylobacteriosis, but more severe disease may also occur.

### Surveillance and control

Campylobacteriosis is notifiable in humans in Norway, but not in animals (except *C. fetus* in cattle). In humans, both campylobacter infections verified by PCR and/or culture are registered in MSIS.

Norway has a surveillance program for *Campylobacter* in broiler chickens. All flocks slaughtered before 51 days of age between the 1<sup>st</sup> May and 31<sup>st</sup> October are tested prior to slaughter. Carcasses from positive flocks are heat treated or frozen prior to sale in order to reduce the potential for transmission to humans. Pasteurisation of milk and disinfection of water are other measures that prevent transmission of *Campylobacter* to humans.

## Results 2019

In MSIS, 4,154 human cases of campylobacteriosis were reported, of which 1,547 contracted the infection in Norway. For 735 of the cases place of infection was unknown. This is an increase compared to the number of cases reported last year and are mainly attributed to a large waterborne outbreak in Norway. Before 2017, cases verified only by PCR were not reported. From 2017, all cases verified by PCR and/or culture are registered in MSIS and reported. However, when the total number of positive cases for 2017, 2018 and 2019 are compared to positive cases verified by both culturing and/or PCR in 2015 and 2016, the numbers for each year are similar, suggesting that there has not been an increase in the occurrence of campylobacteriosis in Norway (Figure 3).

Surveillance in 2019 showed that a total of 103 flocks (5.1%) tested positive for *Campylobacter* spp. before slaughter when all broiler flocks slaughtered before 51 days of age during the period May - October were tested. In total 2,018 flocks from 500 farms were sampled. Of the positive flocks, 30.1% originated from 14 farms (2.8%). A slight decrease was seen in the number of positive flocks and farms compared to 2018.

In the diagnostic services at the NVI, *Campylobacter* was detected in samples from 22 cattle, 20 dogs and one cat. Three wild boar were also *Campylobacter*-positive. For details see the Appendix.




Figure 3. The number of reported cases of campylobacteriosis in humans (data from MSIS) and the percentage of positive broiler flocks (sampled between 1st May and 31st October).



## Evaluation of the current situation

Campylobacteriosis is the most commonly reported zoonosis in humans in Norway. Usually more than half of the cases are reported as infected abroad, but due to a large waterborne outbreak in the western part of Norway, only 45% of the cases were infected abroad in 2019.

Case-control studies have shown that the most common source of campylobacteriosis in Norway is drinking untreated water at home, at holiday homes or in nature. Eating or preparing poultry and barbeque meals have also been identified as risk factors for infection. No studies have demonstrated a link between eating beef or lamb and campylobacteriosis despite a considerable prevalence of *Campylobacter* in these animals in Norway. However, one study showed that eating inadequately heat-treated pork was associated with an increased risk of *Campylobacter* infection. Studies have also shown that direct contact with domestic animals (cattle, sheep, poultry, dogs and cats) is associated with an increased risk of campylobacteriosis in humans.

The prevalence of *Campylobacter* in broilers is low in Norway (3-7% of slaughtered flocks) compared to other countries. The measures implemented in Norway to reduce *Campylobacter* in chicken meat are considered to have had a positive effect on public health. A few farms seem to deliver a high proportion of the positive flocks.

## Yersiniosis

### The disease and its transmission routes

Certain serogroups of the bacteria *Yersinia enterocolitica* can cause disease in humans, for which the most common symptom is diarrhoea. Swine are considered to be the main source of these disease-causing variants. The most common sources of human infection are contaminated food and water.

*Yersinia pseudotuberculosis* is a different bacterium that may cause disease in humans and animals.

### Surveillance and control

Yersiniosis in humans is notifiable, while detection of *Y. enterocolitica* and *Y. pseudotuberculosis* in animals are not. There is no surveillance for this bacterium in animals or food in Norway. Because healthy swine can be carriers, contamination of carcasses may occur at slaughter. Good hygiene at slaughter reduces this risk.

## Results 2019

The number of reported cases of yersiniosis (85) decreased compared to last year, but is still higher than the last few years (Figure 4). All reported cases, but one in 2019 were caused by *Yersinia enterocolitica*. One case was infected with *Y. pseudotuberculosis*.




Figure 4. The number of reported cases of yersiniosis in humans. Data from MSIS.

In the diagnostic services at the NVI, no cases of *Y. enterocolitica* was detected in animals.

## Evaluation of the current situation

Most yersiniosis cases in humans in Norway are sporadic and have been infected domestically. In 2014, there was a significant increase in the number of reported cases due to an outbreak in a military camp and the civilian population. In 2018, two outbreaks were reported, one with six cases due to *Y. enterocolitica* O:3, and one larger outbreak with 20 cases due to *Y. enterocolitica* O:9. Ready to eat salad was suspected to be the source of the second outbreak. There was one outbreak due to *Y. enterocolitica* O:3 in December 2019 in Norway with eleven cases, but it was reported and investigated in January 2020.

*Y. enterocolitica* is presumed to be prevalent in swine and the bacteria cannot be eliminated from swine flocks. During the 1990s routines for improved slaughter hygiene were implemented and this has contributed to reducing the number of human cases of yersiniosis.

## Listeriosis

### The disease and its transmission routes

*Listeria monocytogenes* occurs naturally in the environment and is mainly pathogenic for pregnant women, the elderly and people with a compromised immune system. The main route of infection is in contaminated food or water, and listeriosis can cause fever, abortion, meningitis and septicaemia. In animals, listeriosis also causes meningitis and abortion, and feed is the main source of infection.

### Surveillance and control

Listeriosis in humans is notifiable. In animals, it is categorised as a group C-disease and detection in animals usually does not result in any measures.

Detection of *L. monocytogenes* is part of the control system in the manufacture of certain food products. The upper limit in ready-to-eat foods is 100 cfu/g and 0 cfu/ml in products intended for small children or other vulnerable persons. If the upper limit is exceeded, the food must be withdrawn from market and corrective action must be taken to avoid further contamination. Dietary advice is available; [www.matportalen.no](http://www.matportalen.no) and [www.fhi.no](http://www.fhi.no)

## Results 2019

Twenty-seven cases of listeriosis were reported in humans in 2019 (Figure 5), a slight increase compared to the previous four years. The increase in 2018 and in 2019 can to some extent be attributed to an outbreak of listeriosis in 2018/2019 with 13 cases where the epidemiological and microbiological investigations pointed to “rakfisk”, a fermented fish product, as the cause of the outbreak. As for the rest of Europe an upward trend in the number of listeriosis cases has been observed in the last years.




Figure 5. The number of cases of listeriosis in humans. Data from MSIS.

In the diagnostic services at the NVI, *L. monocytogenes* was detected from a goat herd in connection with abortions, from one sheep and cow, in 2019. The Institute of Marine Research (Havforskningsinstituttet) examined 84 samples of seafood for *L. monocytogenes* and one sample of imported fish products was positive, but had less than 100 cfu/g.

## Evaluation of the current situation

The number of listeriosis cases in both humans and animals in Norway is low, but the infection can have severe consequences. Therefore, it is important that manufacturers of ready-to-eat foods have proper routines in place for preventing *Listeria* in their products, and systems for traceability and withdrawal of products from the market in cases where *L. monocytogenes* are detected. Farmers, especially sheep farmers, must ensure that feed has good quality in order to reduce the risk of listeriosis in animals.

## Verotoxin producing *E. coli* (VTEC)

### The disease and its transmission routes

*Escherichia coli* are normal inhabitants of the intestines of humans and animals. Some *E. coli* can produce verotoxin (also called shigatoxin). These variants are called verotoxin (VTEC) or shigatoxin (STEC) producing *E. coli*, and can cause serious disease and bloody diarrhoea in humans (hence the term EHEC - enterohaemorrhagic *E. coli*). Transmission occur via food, water or by animal contact.

### Surveillance and control

EHEC and diarrhoea-associated haemolytic uremic syndrome (HUS) are notifiable in humans. Detection of VTEC in animals is not notifiable but the NFSA should be informed so that measures can be considered. There is no routine surveillance of VTEC in animals or food, but several screening studies have been performed.

VTEC should not be found in ready-to-eat foods and detection of these bacteria in such foods would lead to withdrawal of the product from the market. Good hygiene and proper routines at slaughter reduces the risk of contamination of meat with VTEC.

## Results 2019

The number of reported EHEC cases in humans (n=512) continues to increase (Figure 6), but the number of cases developing HUS remains low (2-10 cases/year). 51% of the cases in 2019 were diagnosed with low-virulent VTEC.




Figure 6. The number of reported cases of EHEC (enterohaemorrhagic *E. coli*) in humans. Data from MSIS.

No food samples were received for analyses at the Norwegian Veterinary Institute in relation to investigation of human EHEC infections in 2019.

## Evaluation of the current situation

The occurrence of EHEC-infections reported in humans has been increasing since 2012. More than half of the cases have been infected in Norway. The increase is most likely associated with the introduction of culture independent diagnostics (PCR) as a routine in primary diagnostics and the fact that more patients than before are investigated for EHEC. Several major medical microbiological laboratories investigate all submitted faecal specimens for several different pathogens, including EHEC. Previously, analysis for EHEC was only performed based on defined clinical or epidemiological indications.

In a survey of zoonotic *E. coli* in Norwegian cattle, conducted in 2014, the Norwegian Veterinary Institute found a low occurrence of STEC, with 15.6% of 179 herds positive for at least one STEC belonging to six serogroups (O26, O91, O103, O121, O145 and O157).

## Tuberculosis

### The disease and its transmission routes

Tuberculosis is caused by species in the *Mycobacterium tuberculosis*-complex. As a zoonosis, *Mycobacterium tuberculosis* subsp. *bovis* (*M. bovis*), which causes bovine tuberculosis, is the most important. This bacterium is mostly found in cattle. Humans are usually infected by drinking unpasteurised milk. Tuberculosis in humans is usually caused by *M. tuberculosis* subsp. *tuberculosis* (*M. tuberculosis*) which is transmitted between humans in microscopic airborne droplets. Humans may also transmit tuberculosis to animals. Tuberculosis can cause an array of symptoms depending on the affected organ system, but symptoms from the respiratory system are most common. Tuberculosis is a chronic infection in both animals and humans.

### Surveillance and control

Tuberculosis in humans is notifiable in Norway. Persons in higher-risk groups are offered BCG vaccination. Tuberculosis caused by *M. bovis* and *M. tuberculosis* in animals is categorised as a group B disease, while detection of other non-tuberculous mycobacteria are group C.

Norway is free of bovine tuberculosis, and this is acknowledged in the EEA agreement where Norway is declared as officially free. Vaccination of animals against tuberculosis is forbidden in Norway. All animals, except poultry, are inspected for tuberculosis at commercial slaughter. Any suspicious findings will be examined further. Tuberculin testing is performed on all breeding bulls and breeding boars at semen collection facilities, imported animals, and in cases where tuberculosis is suspected or must be excluded. Animals with a positive tuberculin test will be euthanized and further examined. The NFSA have a surveillance program for *M. tuberculosis* in cattle, camelids and farmed deer.

## Results 2019

In total, 175 cases of tuberculosis in humans were reported in 2019. One of these was caused by *M. bovis*, and the infection was contracted abroad.

All cattle, sheep, goats, swine and horses commercially slaughtered were examined *post mortem*. In addition, 205 breeding pigs and 254 breeding bulls were tuberculin tested, and all were negative. As part of diagnostic testing, samples from 24 swine, four alpaca and one llama were tested for *Mycobacterium* spp. *Mycobacterium bovis* was not detected in any of the samples.

## Evaluation of the current situation

*M. bovis* infection in humans is rarely reported in Norway. Less than 1% of the reported human tuberculosis cases in the last 10 years were caused by *M. bovis*, and these patients were either infected abroad or many decades ago in Norway. Since the mid-1990s, the number of tuberculosis cases caused by *M. tuberculosis* has increased in Norway due to immigration, but for the last 4 years (since 2013) the number has decreased.

Bovine tuberculosis, *M. bovis* infection in cattle, was eradicated in Norway in 1963, but was detected in one area in the 1980s. This was most probably transmission from an infected human. Tuberculosis in animals caused by *M. tuberculosis* is rare in Norway and was last reported in a dog in 1989.

Import of live animals to Norway, especially camelids like llama and alpaca, is associated with a risk of introducing *M. bovis* to the Norwegian animal population. Foreign farm labourers could represent a potential, but low risk of introducing *M. bovis* and *M. tuberculosis* to Norwegian animals.

## Brucellosis

### The disease and its transmission routes

Brucellosis is caused by *Brucella* bacteria, of which *B. abortus* (cattle), *B. melitensis* (sheep), and *B. suis* (pigs) are the most important zoonotic species. *B. canis*, which causes disease in dogs, is less pathogenic for humans.

Brucellosis may cause sterility and abortion in animals. In humans, fever is the most common symptom. The bacteria are shed in milk, and humans are usually infected through consumption of unpasteurised milk and products made from unpasteurised milk.

### Surveillance and control

Brucellosis in humans is notifiable and brucellosis in animals is listed as a notifiable group A-disease.

The surveillance program for *Brucella* includes blood tests from cattle that have aborted and annual blood testing of a sample of the sheep and goat population. In addition, breeding bulls and boars and imported animals are tested. Vaccination of animals against brucellosis is forbidden in Norway. Norway is official free of brucellosis in cattle according to the EEA agreement. Although not officially recognized as such, Norway is also free from brucellosis in the small ruminants and swine populations.

## Results 2019

Four cases of brucellosis in humans were reported in 2019, three were infected abroad and for one case the place of infection was unknown.

As part of the surveillance programmes, 104 cattle from 31 herds 8,950 sheep from 33,363 flocks, and 1,751 goats from 58 herds were tested. Antibodies against *Brucella* spp. were not detected. In addition, 323 breeding cattle and 2049 breeding pigs were tested and were negative for antibodies against *Brucella* spp. Eight alpaca were tested and were negative after import to Norway. Five dogs tested negative for *B. canis* antibodies and/or by culture.

## Evaluation of the current situation

In humans, brucellosis is rare with only 0-4 reported cases per year, most of which have been infected abroad. Some have been infected domestically from laboratory work or from eating products purchased abroad that were made from unpasteurised milk.

Bovine brucellosis was eradicated from Norway in 1953 and brucellosis in sheep, goats and pigs has never been detected in Norway. *B. canis* has been detected in Sweden, but not in Norway.

## Trichinellosis

### The disease and its transmission routes

Trichinellosis is caused by small round worms, called *Trichinella*. Animals and humans may be infected through consumption of raw or poorly heat treated meat containing larvae. In the intestines, the larvae grow into adult worms and reproduce. Adult females set free larvae that move away from the intestines to muscle tissue. The most common symptom of trichinellosis is muscle pain, but the disease can also take more serious forms. Raw or poorly heat treated meat is the main source of infection.

### Surveillance and control

Trichinellosis in humans is notifiable, and in animals it is a group B-disease. All carcasses of pigs and horses are checked for the presence of *Trichinella* at slaughter. Positive carcasses will be destroyed. Predator animals that are hunted/slaughtered and used for consumption (eg. wild boar or bear) should also be tested for *Trichinella*.

## Results 2019

No cases of trichinellosis were reported in humans in 2019.

All commercially slaughtered pigs and horses were tested for *Trichinella*, and none were positive. The NVI also recommend that all hunted wild boar are tested for *Trichinella* before consumption. In the health surveillance program for wild boar, which started in 2018, samples from 79 wild boar harvested through hunting were tested for *Trichinella* in 2019, none were positive.

## Evaluation of the current situation

Trichinellosis in humans is very rare in Norway. The last case was reported in 1996, and the last case infected in Norway was reported in 1980.

*Trichinella* in domestic animals in Norway was last reported in two pig herds in 1994, and before that in 1981. *Trichinella* may be found in wild animals, and the parasite may transmit to domestic animals kept outside such as swine and horses. In 2018, 15 hunted wild boar were tested for *Trichinella* and all were negative.

## Echinococcosis

### The disease and its transmission routes

*Echinococcus granulosus* and *E. multilocularis* are small tape worms that can cause serious disease in humans. The parasites have their adult stage in the intestines of predators (eg. fox and dog), and parasite eggs are shed in faeces of these hosts (definitive host). Other animals (intermediate host) are infected through ingestion of the eggs. In the intermediate host the eggs hatch to larvae that migrate and encapsulate in cysts in various organs. The intermediate host must be eaten by a definitive host for the parasite to develop further into adult stages. It is the larval cysts in the intermediate host, e.g. in humans, that cause disease. Humans may be infected through eating fruit and berries contaminated with eggs or through direct contact with infective definitive hosts (e.g. dogs).

### Surveillance and control

Echinococcosis in humans is notifiable in Norway and in animals it is a group B disease. Intermediate hosts for *E. granulosus* (eg. reindeer and cattle), are examined at slaughter. Since 2006, hunted red foxes have also been examined for *E. multilocularis*. This surveillance was intensified in 2011 when the parasite was detected in Sweden.

Dogs entering Norway must be treated for *Echinococcus* before arrival. Regular anti-parasitic treatment of dogs is also recommended in areas with reindeer.

## Results 2019

Seven cases of echinococcosis in humans were reported in 2019, and all had contracted the infection abroad.

In the surveillance program for *E. multilocularis*, 541 foxes and 19 wolves were examined, and *E. multilocularis* was not detected in any of them. All commercially slaughtered cattle, sheep and pigs were examined for *Echinococcus post mortem*, and no cases were identified. For details see the Appendix.

## Evaluation of the current situation

Echinococcosis has never been a public health problem in Norway. In humans between 0 and 7 cases have been reported annually of which all cases have been infected abroad.

*E. granulosus* was common in reindeer in northern Norway until the 1950s. Systematic treatment of shepherd dogs and reduced feeding of these dogs with raw meat and offal was effective and the parasite is now very rare in reindeer. It was last detected in 1990 and 2003. In cattle, *E. granulosus* was last reported in 1987.

*E. multilocularis* has never been detected in mainland Norway. However, it is detected in Sweden, and surveillance of red foxes has been intensified in Norway in order to rapidly detect the parasite should it be introduced to Norway. Since 2002, 5,037 red foxes have been tested, and all were negative.

*E. multilocularis* is endemic in Svalbard in sibling vole (*Microtus levis*) and the Arctic fox (*Vulpes lagopus*). Dogs and people in Svalbard are therefore at risk.

It is essential that dog owners follow regulations on antiparasitic treatment when entering Norway from abroad. Echinococcosis occurs in dogs in southern Europe, and the infection may be introduced to the Norwegian population of intermediate and definitive hosts via untreated, imported dogs or dogs returning with their owners after holidays abroad.



## Toxoplasmosis

### The disease and its transmission routes

*Toxoplasma gondii* is a single celled parasite that has its adult stage in the cat (definitive host). The parasite is shed in faeces and intermediate hosts (e.g. sheep, human, rodents) are infected through contaminated food or water or by direct contact with contagious cats. Humans can also be infected through consumption of inadequately heat treated meat. Healthy adults will usually not become sick from toxoplasmosis. However, if women contract the infection for the first time during pregnancy, it may result in abortion or harm the foetus.

### Surveillance and control

Toxoplasmosis is not notifiable in humans or animals in Norway.

The NFSA provides dietary advice to persons in risk groups ([www.matportalen.no](http://www.matportalen.no)). Every year some animals are tested for *T. gondii* due to disease, abortion or in association with import/export. Testing of cats for *T. gondii* is not considered necessary.

## Results 2019

As part of the diagnostic work at the NVI, 11 sheep were tested serologically for *T. gondii*. None were positive.

## Evaluation of the current situation

*T. gondii* is prevalent in Norway, but is less prevalent than in southern Europe. It has been estimated that 90% of Norwegian women are susceptible to infection, and that 2 in 1,000 pregnant women contract the infection for the first time during pregnancy. The parasite is estimated to transmit to the foetus in approximately 50% of these cases.

*T. gondii* is prevalent in several mammals in Norway, in particularly cats and sheep. In an investigation of lambs in the 1990s, 18% of the tested lambs had antibodies against *Toxoplasma*, and positive animals were found in 44% of the tested flocks. Similarly, in a study performed between 2002 and 2008, 17% of tested goats were antibody-positive, and positive animals were found in 75% of the tested herds. In another study, performed in the 1990s, 2.6% of pigs for slaughter were antibody positive.

Wild deer may be infected with *T. gondii*. In a serological study of 4,300 deer hunted between 1992 and 2000, 34% roe deer, 13% elk, 5% hart deer and 1% reindeer were antibody positive.

## Rabies

### The disease and its transmission routes

Rabies is caused by a lyssavirus, and the infection manifests itself as a neurological disease. The virus transmits through bites, or from exposure of open wounds to saliva from rabid animals. The incubation period is usually 1-3 months but may be longer. Untreated rabies is fatal. In Europe, classic rabies and bat rabies are caused by different viruses. Bat rabies in Europe has a much lower zoonotic potential than classic rabies.

### Surveillance and control

Rabies is notifiable both in humans and in animals (group A disease). A vaccine is available for people who are traveling to high risk areas for extended periods. The vaccine is also used in combination with anti-serum to treat people who may have been exposed to rabies.

Animals with rabies will be euthanized, and measures will be implemented to stop further spread. From the 1<sup>st</sup> January 2012, dogs and cats imported from EU and EEA countries are only required to be vaccinated against rabies. Previously, a blood test to prove sufficient antibody titres was also mandatory. For dogs and cats imported from non EU non EEA countries, both a rabies blood test and proof of antibody titre is required.

## Results 2019

Rabies was detected in one person in Norway in 2019. The disease was contracted abroad.

Two dogs and five bats from mainland Norway, and three arctic foxes and six Svalbard reindeer (from Svalbard), were tested for rabies at the NVI. Rabies was not detected in any of these. For further information see the Appendix.

## Evaluation of the current situation

In rare cases, bat rabies may transmit from bats to other warm-blooded animals, including humans. Therefore, care is advised when handling bats, and any bite from a bat should be consulted with a doctor. It is not considered necessary to start vaccinating animals in Norway due to the detection of bat rabies in 2015.

Classic rabies has never been detected in mainland Norway, but it has been detected in Arctic fox, reindeer and a ringed seal in Svalbard. The last detection was in 2011-2012 and before that 1999. Hence, outbreaks of rabies occur sporadically in Svalbard, most probably due to migrating arctic foxes during winter. It is important that persons living in or traveling to Svalbard are aware that rabies may occur among wild animals and take necessary precautions.

Dogs imported to Norway without vaccination may confer a risk of introducing rabies to mainland Norway. In a study performed at the NVI in 2012, serological results indicated that approximately 50% of dogs imported from Eastern Europe were improperly vaccinated or not vaccinated at all. Illegal import of dogs to Norway poses a threat to human and animal health due to the risk of introducing rabies to the country.

## Q-fever

### The disease and its transmission routes

Q-fever is caused by the bacteria *Coxiella burnetii*, and is mainly associated with ruminants. However, humans and other animals may also become infected and sick. The bacteria are shed in urine, faeces, foetal fluids, placenta and foetal membranes, and can survive for extended periods in the environment. Transmission is airborne via aerosols. In animals, infection results in weak offspring, abortions, infections of the placenta and uterus. In humans *C. burnetii* may cause influenza-like symptoms and rarely more serious disease.

### Surveillance and control

Q-fever in humans has been notifiable in Norway since 2012, and is a group C-disease in animals. Animals with clinical signs of Q-fever must not have contact with animals from other herds/farms and the NFSA may impose restrictions on animal holdings where infection is confirmed or suspected.

From 2012, samples collected in the surveillance programme for *Brucella abortus* in cattle have also been tested serologically for *C. burnetii*. The programme involves passive clinical surveillance, and blood samples from cattle with an abortion in the second half of the pregnancy are analysed.

## Results 2019

Eight cases of Q-fever in humans were reported in 2019. Seven had contracted the infection abroad and for one case the place of infection was unknown.

At the NVI, blood samples from a total of 2,226 cattle, 2,056 sheep and 208 goats were tested serologically for *C. burnetii* in the surveillance program. In addition 17 cattle, 18 sheep and 2 alpaca were tested in relation to export/import. Antibodies were not detected in any samples. For further information see the Appendix.

## Evaluation of the current situation

Q-fever is not currently a problem for human or animal health in Norway. The infection became notifiable in humans in 2012, and since then 25 cases have been reported in total. Of these, 23 cases were infected abroad and one case had an unknown place of infection.

Q-fever has not been detected in Norwegian animals. Screening studies were performed in 2008 (460 bovine dairy herds and 55 bovine meat herds), in 2009 (349 goat herds and 45 bovine herds) and in 2010 (3,289 bovine dairy herds). Since then, testing has been performed on imported animals and as part of diagnostic testing of sick animals.

## BSE and vCJD

### The disease and its transmission routes

Bovine spongiform encephalopathy (BSE, mad cow disease) in cattle and Creutzfeldt-Jacob disease (CJD) in humans are transmissible spongiform encephalopathies (TSE). These fatal diseases cause spongy degeneration of the brain and spinal cord. The infective agents are prions, protein structures without DNA. A form of CJD, variant CJD (vCJD) was first described as the cause of death in a person in the UK in 1995. The disease was suspected to be caused by consumption of beef containing the prion associated with classic BSE.

Other TSE-diseases that do not transmit between animals and humans have also been described, such as atypical BSE in cattle, scrapie in sheep, sporadic CJD in humans and chronic wasting disease (CWD) in deer.

### Surveillance and control

Surveillance for BSE started in Norway in 1998, and includes testing of imported animals and their offspring, emergency slaughtered cattle, cattle with defined clinical signs at slaughter and a sample of routinely slaughtered cattle. All small ruminants with scrapie are tested to rule out BSE.

At slaughter, specified risk material (SRM) is removed from cattle and small ruminants. It is forbidden to use protein from animal (including fish protein) in feed for ruminants. Norway banned the use of bone meal in ruminant feed in 1990.

## Results 2019

No cases of sporadic vCJD were reported in humans in 2019.

In total, 6,884 cattle were tested, and all were negative for BSE.

## Evaluation of the current situation

The situation with respect to classic BSE has always been favourable in Norway, largely due to restricted and controlled import of live animals, meat and bone meal in the past when the disease emerged and spread in Europe, and historical strict regulations on heat treatment and use of meat and bone meal.

The first and only case of BSE in cattle in Norway, an atypical BSE case - and as such not a zoonosis, was detected in 2015.

## Antimicrobial resistance

Antimicrobial resistant bacteria may be zoonotic and transmit through direct or indirect contact, including through food. One example is methicillin resistant *Staphylococcus aureus* (MRSA), which may transmit between animals and humans.

### Surveillance and control

Infection and carriage of MRSA in humans is notifiable in Norway. In addition, selected microbes from certain infections, and their resistance profiles, are reported annually to the NORM surveillance programme for antimicrobial resistance in human pathogens.

In 2000, Norway implemented a surveillance programme for antimicrobial resistance in pathogens from animals, feed and food (NORM-VET). In 2013, a separate surveillance program for MRSA in swine was established. Norway has chosen a strategy to eradicate MRSA from swine, and therefore detection of MRSA in any production animal is reported to the NFSA. Detection of MRSA and several other forms of resistant bacteria in animals became notifiable in Norway in 2019.

## Results 2019

Details on detection of selected pathogens in humans and animals and their antimicrobial resistance profiles are presented in the annual NORM/NORM-VET report (<https://www.vetinst.no/overvaking/antibiotikaresistens-norm-vet>).

The surveillance programme in 2019 detected one pig herd with MRSA. In total, 722 herds were included in the survey, of which 79 were genetic nucleus or multiplier herds, 12 were central units of the sow pool, 22 were farrow to grower or farrow to finish herds, and the remaining 609 were herds with more than 10 sows. Additional samples from six farrow to finish herds were received, though not included in the 2019 surveillance. MRSA was detected from one of these. Contact tracings from the two positive pig herds were conducted, resulting in an additional 6 MRSA positive herds. Eradication was imposed in all eight herds.

In 2019, a total of seven *Salmonella* spp. isolates from animals were susceptibility tested. The seven isolates included one each from turkey, poultry, dog and pig, and three from cats. The isolates were fully susceptible to the antimicrobial agents tested for.

*Campylobacter coli* from fattening pigs were retrieved from samples taken at the abattoirs throughout the year. Overall, 46.6% of the isolates were resistant to at least one antimicrobial agent, but none of the isolates were multidrug resistant (i.e. resistant to  $\geq$  three antimicrobial classes).

Nine pathogenic *Y. enterocolitica* isolates were obtained from a survey of minced pork meat. All nine isolates were fully susceptible to all antimicrobial agents included in the test panel.

## Evaluation of the current situation

Antimicrobial resistance in bacteria has become a serious threat to human and animal health globally. The prevalence of antibiotic resistant bacteria is still low in both humans and animals in Norway compared to other European countries. However, the situation is under threat by the high use of antibiotics globally.

Resistant pathogens may spread through healthy carriers. MRSA was most likely first introduced to Norwegian swine production through foreign labourers carrying the bacteria, and subsequently spread further through movement of live animals. From swine, MRSA may transmit back to humans through direct or indirect contact. This form of transmission is difficult to control, and in this respect MRSA is an example of a modern biosecurity challenge in Norwegian food production.

## Foodborne outbreaks

An outbreak is either defined as more cases than expected of a specific disease within a defined geographical area and time period, or as two or more cases of a disease with a common source of infection. In 2005, the NIPH and the NFSA introduced a web-based system for reporting outbreaks, Vesuv. The system is used by specialist- and municipal health services and the NFSA to notify outbreaks. The following types of outbreaks are notifiable through Vesuv: outbreaks of conditions that are notifiable in MSIS; outbreaks associated with food or water; outbreaks caused by particularly serious infections; very large outbreaks; and outbreaks in healthcare institutions. The four last categories also include outbreaks of conditions that are not notifiable in MSIS.

The purpose of investigating foodborne outbreaks is to stop the outbreak, implement control measures and prevent future outbreaks. The District Medical Officer is responsible for coordinating investigation and response to outbreaks in his/her municipality. Proper outbreak investigation requires cooperation between local and central health authorities, the NFSA and other relevant authorities.

### Results 2019

In 2019, the NIPH received 46 notifications through Vesuv of possible or confirmed foodborne outbreaks outside health institutions. In total, 2, 705 persons were reported to have become sick in these outbreaks. The number of affected persons in each of the outbreaks varied between 2 and 2000 (median 12) The most common infective agent was Norovirus (13 outbreaks) followed by enterohaemorrhagic *E. coli* (EHEC) and *Campylobacter* (5 outbreaks each). In 8 of the outbreaks, the causative agent was not identified (Figure 7).




Figure.7. The number of reported foodborne outbreaks where an agent was verified or strongly suspected.

## Appendix Tables 2019

Table 1. Human population of Norway

Table 2. Animal population of Norway

Table 3. *Salmonella* serovars in humans in Norway

Table 4. Human cases of campylobacteriosis distributed by county

Table 5. Foodborne outbreaks

Table 6. *Salmonella* in feed and feedstuff

Table 7. *Salmonella* in animals

Table 8. *Salmonella* in food

Table 9. Selected zoonoses in animals

Table 1. Human population of Norway per 1st January 2019 (from statistics Norway).

Age group	Female	Male	Total
0 - 9	298 981	315 935	614 916
10 - 19	311 315	329 226	640 541
20 - 29	346 473	365 946	712 419
30 - 39	349 639	367 730	717 369
40 - 49	354 815	373 824	728 639
50 - 59	337 657	354 087	691 744
60 - 69	288 032	288 715	576 747
70 - 79	217 710	202 128	419 838
80 - 89	106 722	74 309	181 031
90 -	298 981	13 173	44 968
Total	2 643 139	2 685 073	5 328 212

Table 2. Animal population of Norway in 2019.

Animal species - category	Number*		
	Herds/flocks	Animals	Slaughtered animals
Cattle - total	13 500 <sup>a</sup>	862 000 <sup>a</sup>	304 400 <sup>c</sup>
Dairy production	6 500 <sup>a</sup>	177 000 <sup>a</sup>	
Meat production	4 500 <sup>a</sup>	79 600 <sup>a</sup>	
Other stock	880 <sup>a</sup>	42 000 <sup>a</sup>	
Sheep - total	14 600 <sup>a</sup>	2 469 000 <sup>a</sup>	1 178 000 <sup>c</sup>
Sheep >1 year	14 000 <sup>a</sup>	937 000 <sup>a</sup>	
Goats - total	1 300 <sup>a</sup>	71 000 <sup>a</sup>	27 700 <sup>c</sup>
Dairy goats	350 <sup>a</sup>	36 300 <sup>a</sup>	
Swine - total	2 000 <sup>a</sup>	761 000 <sup>a</sup>	1 622 000 <sup>c</sup>
Breeding pigs	1 100 <sup>a</sup>	45 200 <sup>a</sup>	
Slaughter pigs	1 800 <sup>a</sup>	420 000 <sup>a</sup>	
Chickens ( <i>Gallus gallus</i> )			
Grandparent stock - egg producers	2 (2) <sup>b1</sup>		
Parent stock - egg producers	7 (23) <sup>b1</sup>		
Parent stock - broiler	94 (210) <sup>b1</sup>		
Laying hens	592 (874) <sup>b</sup>		
Broilers	586 (4 511) <sup>b</sup>		68 288 000 <sup>c</sup>
Turkey, goose and duck			
Parent stock	7 (17) <sup>b1</sup>		
Meat production	51 (250) <sup>b</sup>		822 700 <sup>c</sup>
Horse			108 <sup>c</sup>
Reared deer	80 <sup>d</sup>	7 500 <sup>d</sup>	

<sup>a</sup> Figures from the registry of production subsidy per 31.3.2019.

<sup>b</sup> Figures from the surveillance programme for Salmonella

<sup>c</sup> Figures from the Norwegian Agriculture Agency (based on delivery for slaughter)

<sup>d</sup> Figures from the Norwegian Agriculture Agency - per 31.7.2019

<sup>1</sup> Production flocks only.



Table 3. The most common *Salmonella* serovars found in humans in Norway in 2019.

Serovar	Place of infection			Total
	Norway	Abroad	Unknown	
<i>S. Enteritidis</i>	73	280	45	398
<i>S. Typhimurium</i>	47	46	10	103
<i>S. Typhimurium monophasic variant</i>	19	29	11	59
<i>S. Stanley</i>	17	28	6	51
<i>S. Agbeni</i>	56	0	0	56
<i>S. Java</i>	4	16	3	23
<i>S. Newport</i>	5	14	3	22
<i>S. Coeln</i>	11	2	1	14
<i>S. Infantis</i>	3	8	1	12
<i>S. Agona</i>	0	10	2	12
<i>S. Napoli</i>	5	4	1	10
<i>S. Chester</i>	2	8	0	10
<i>S. Saintpaul</i>	2	7	0	9
Others	73	194	48	315
Total	317	646	131	1094

Table 4. Human cases of campylobacteriosis (infected in Norway) in 2019 distributed by county. From 2017 both cases verified by PCR and/or culturing are notifiable to MSIS and included in the table. PCR positive cases are also included for 2015 and 2016.

County	2011	2012	2013	2014	2015	2016	2017	2018	2019
Akershus	91	108	120	157	123	142	127	94	126
Aust-Agder	18	20	14	33	14	23	29	12	17
Buskerud	48	63	60	65	63	68	83	56	84
Finnmark	13	6	6	19	12	21	19	21	18
Hedmark	39	28	51	54	42	58	56	32	47
Hordaland	136	128	115	156	122	148	185	163	338
Møre og Romsdal	54	36	47	73	34	35	71	59	47
Nordland	47	31	46	60	47	42	44	57	45
Nord-Trøndelag	28	31	29	34	27	42	33		
Oppland	69	68	62	76	57	66	96	81	66
Oslo (f)	113	136	103	154	120	109	109	144	125
Rogaland	177	124	169	130	124	166	205	129	117
Sogn og Fjordane	45	24	27	27	26	20	31	25	12
Sør-Trøndelag	85	115	95	117	94	98	116		
Telemark	37	39	34	50	35	55	46	32	46
Troms	25	20	23	55	43	57	52	35	47
Trøndelag								118	130
Vest-Agder	45	34	41	54	33	40	48	32	34
Vestfold	42	51	42	79	48	186	101	76	192
Østfold	63	52	66	71	44	73	61	49	54
Total	1 175	1 114	1 150	1 464	1 108	1 449	1 512	1 215	1 545

Table 5. Foodborne outbreaks 2019.

Agent	2011	2012	2013	2014	2015	2016	2017	2018	2019
<i>Salmonella</i> sp.	1	4	2	1	1		2	4	1
<i>Campylobacter</i> sp.	5	2	2	3	1	2	3	1	7
<i>Yersinia</i> sp.	1		1	2	1	1	1	2	
<i>Listeria monocytogenes</i>		1	1					1	1
<i>Escherichia coli</i> (VTEC)		1	2	1		1	3	4	6
<i>Clostridium</i> sp.	3	2			1		1		1
<i>Bacillus cereus</i>	1	1		4	4				
<i>Staphylococcus enterotoxin</i>	2				3		1		1
Parasites			1			2	2	1	2
Virus	14	13	15	14	13	11	7	17	16
Other	8	2	2	2	1	1	1	5	3
Unknown	14	18	36	28	15	11	15	17	8
Total	49	44	62	55	40	29	36	52	46

Table 6. *Salmonella* in feed and feedstuff 2019.

Category	Number tested*	Number positive	Comment
Feedstuff			
Cereal grain	131	1	<i>S. Cubana</i>
Corn	58	0	
Rape	73 (1)	4	<i>S. Agona</i> (2), <i>S. Kentucky</i> , <i>S. Typhimurium</i>
Soya	2 498 (4)	0	
Sunflower	25	0	
Legume seeds etc.	30	0	
Tubers, roots etc.	38	0	
Other plant based feedstuffs	32 (1)	1	<i>S. Putten</i>
Meat based feedstuff	322	3	<i>S. Rissen</i> , <i>S. Havana</i> (2)
Marine based feedstuff	126	0	
Feed			
Domestic animals (cattle, swine, poultry)	57 (76)	0	
Fish	2 842	3	<i>S. Bradford</i> , <i>S. Livingstone</i> (2)
Fur animals	100	0	
Environmental samples in factories producing feed and feedstuff	9 903	190	31 different serovars

\* Total numbers are presented, in brackets the number of samples collected by Authorities.

Table 7. *Salmonella* in animals 2019.

Category	Number* tested	Number* positive	Comment
Chicken - surveillance - breeding flocks	233	0	
Chicken - surveillance - layer flocks	874	0	
Chickens - surveillance - broiler flocks	4 511	1	<i>S. Give</i>
Chicken flocks - other samples	12	0	
Turkey, ducks, geese - surveillance - breeding flocks	17	0	
Turkey, ducks, geese - surveillance - meat flocks	250	1	<i>S. Agona</i>
Turkey, ducks, geese - other samples	2	0	
Cattle - surveillance - animals	3 151	0	
Cattle - diagnostics - herds	35	0	
Sheep - diagnostics - herds	37	5	<i>S. diarizonae</i>
Goats- diagnostics - herds	2	0	
Swine - surveillance - slaughter pigs - animals	1 774	2	<i>S. Hessarek</i> , <i>S. Typhimurium</i>
Swine - surveillance - sows - animals	1 645	0	
Swine - surveillance - breeding herds	77	0	
Swine - diagnostics - herds	7	0	
Horse - diagnostics - animal	224	2	<i>S. Typhimurium</i> monophasic
Dogs - diagnostics	156	1	<i>S. Typhimurium</i>
Cat - diagnostics	21	2	<i>S. Typhimurium</i> , <i>S. Enteritidis</i>
Alpaca- herds - diagnostics	2	0	
Camel- herds - diagnostics	1	0	
Wild boar - surveillance - animals	71	0	
Animals/birds/zoo birds/zoos	24	0	
Reptiles**	19	16	<i>S. Carmel</i> (1), <i>S. diarizonae</i> (9), <i>S. houtenae</i> (1), <i>S. arizonae</i> (1), <i>S. Havana</i> (1), <i>S. Kisarawe</i> (1), <i>S. Tennessee</i> (1), <i>Salmonella</i> sp.(1)
Various wild animals	11	0	
Wild birds	12	0	

\* Units for numbers are given in the first column.

Table 8. *Salmonella* in food 2019.

Category	Number sampled	Number positive	Comment
Cattle - swab of carcass - surveillance	3 122	0	
Swine - swab of carcass - surveillance	3 314	0	
Meat scrapings (cattle, swine. sheep) - surveillance	3 082	2	<i>S. diarizonae</i>
Fish - Norwegian - IMR*	20	0	
Fish - Imported - IMR*	94	0	
Shellfish - Norwegian - IMR*	26	0	

\* Institute of Marine Research (Havforskningsinstituttet)

Table 9. Selected zoonoses in animals 2019. *Salmonella* is presented in separate tables.

Infection/agent	Category	Number tested	Number positive	Comment
Campylobacteriosis	Broiler chicken flocks - surveillance	2 108	103	May - October
	Cattle - diagnostics	52	22	<i>C. jejuni</i> (20), <i>C. sp.</i> (2)
	Goat - diagnostics	1	0	
	Wild boar - surveillance	63	3	<i>Campylobacter sp.</i>
	Dog - diagnostics	118	20	<i>C. upsaliensis</i> (14), <i>C. jejuni</i> (6)
	Cat - diagnostics	1	1	<i>C. upsaliensis</i>
Tuberculosis	Cattle - tuberculin testing	254	0	
	Swine - tuberculin testing	205	0	
	Swine - diagnostics	24	15	<i>M. avium</i> subsp. <i>hominissuis</i>
	Alpaca - surveillance	4	0	
	Lama - surveillance	1	0	
Brucellosis	Cattle - surveillance	104	0	
	Cattle - breeding animals, export	323	0	
	Sheep - surveillance	8 950	0	
	Goat - surveillance	1 751	0	
	Swine - breeding stock	2 049	0	
	Dog	5	0	
	Alpaca - import	8	0	
Echinococcosis	Fox - surveillance	541	0	
	Wolf - surveillance	18	0	
	Cattle, small ruminants, swine, horse	All slaughtered*	0	
Toxoplasmosis	Sheep - diagnostics	11	4	
Rabies	Dog - diagnostics	2	0	
	Arctic fox - diagnostic	3	0	
	Svalbard reindeer - diagnostic	6	0	
	Bats - diagnostic	5	0	
Trichinellosis	Pig and horse	All slaughtered*	0	
Q-fever	Cattle - surveillance	2 226	0	
	Cattle - export	17	0	
	Sheep - surveillance	2056	0	
	Sheep - import	18	0	
	Goat - surveillance	208	0	
	Alpaca - import	2	0	
BSE	Cattle	6 884	0	
MRSA	Swine - surveillance	722	1	CC398 t034

\* Commercial slaughter (for animal population see Table 2.).


*Scientifically ambitious, forward-looking and cooperatively oriented  
– for integrated health*

The Norwegian Veterinary Institute is a national research institute that operates in the fields of animal and fish health, food safety and feed hygiene; its primary task is to provide the authorities with independently generated knowledge.

Emergency preparedness, diagnostic services, monitoring, reference functions, consulting, and risk assessments are all important areas of activity. Our products and services include research results and reports, analyses and diagnoses, studies and advice.

The Norwegian Veterinary Institute's central laboratory and administration lie in Oslo, and we operate regional laboratories in Sandnes, Bergen, Trondheim, Harstad and Tromsø.

The Norwegian Veterinary Institute collaborates with a large number of national and international institutions.



### Fish health



### Animal health



### Food safety



**Oslo**  
postmottak@vetinst.no

**Trondheim**  
vit@vetinst.no

**Sandnes**  
vis@vetinst.no

**Bergen**  
post.vib@vetinst.no

**Harstad**  
vih@vetinst.no

**Tromsø**  
vitr@vetinst.no

[www.vetinst.no](http://www.vetinst.no)



**Veterinærinstituttet**  
Norwegian Veterinary Institute