

The surveillance programme for avian influenza (AI) in poultry in Norway 2014

Bruce David

Siri Kulberg Sjurseth

Chiek Er



Surveillance programmes for terrestrial and aquatic animals in Norway

Annual report 2014

Project managers at the Norwegian Veterinary Institute:

Ståle Sviland (Terrestrial animals)

Anne-Gerd Gjevre (Aquatic animals)

Mona Torp (Food safety)

Publisher

Norwegian Veterinary Institute

PO Box 750 Sentrum

N-0106 Oslo

Norway

Fax: + 47 23 21 60 95

Tel: + 47 23 21 60 00

E-mail: postmottak@vetinst.no

www.vetinst.no

ISSN 1894-5678

Title:

The surveillance programme for avian influenza (AI) in poultry in Norway 2013

Authors:

Bruce David, Siri Kulberg Sjurseth, Chiek Er

Date: 2015-03-24

Front page photo: Processed from Colourbox

Any use of the present data should include specific reference to this report.

Example of citation:

David B, Sjurseth SK, Er C. The surveillance programme for avian influenza (AI) in poultry in Norway 2014. *Surveillance programmes for terrestrial and aquatic animals in Norway. Annual report 2014*. Oslo: Norwegian Veterinary Institute 2015.

The surveillance programme for avian influenza (AI) in poultry in Norway 2014

Bruce David, Siri Kulberg Sjurseth, Chiek Er

Surveillance in 2014 did not detect infection with highly pathogenic influenza A virus in poultry.

Introduction

The Norwegian Food Safety Authority is responsible for implementing the surveillance programme for avian influenza (AI) in poultry. The programme, which was started in 2006, is based on serological investigations of poultry. The Norwegian Veterinary Institute is responsible for planning, laboratory investigations and reporting components of the programmes.

AI is a serious, highly contagious disease of poultry and other captive birds caused by many different subtypes of influenza type A viruses. The level of risks posed by the different subtypes for animal and public health is very variable due to rapid virus mutation and possible re-assortment of the genetic material between different subtypes.

Current knowledge indicates that the health risks posed by the so-called Low Pathogenic AI (LPAI) viruses are lower than that posed by Highly Pathogenic AI (HPAI) viruses. The HPAI viruses originate from a mutation of LPAI viruses of either H5 or H7 subtype. HPAI can cause disease in poultry resulting in mortality rate exceeding 90 %.

In general, domestic poultry populations are free from AI viruses. However, wild waterfowls are the natural reservoirs for all influenza A virus subtypes. Infected birds do not usually develop clinical disease, but may shed large amounts of virus upon infection (1). The national surveillance programme for AI in poultry was started in 2006 and modelled on EU's Council Directive 2005/94/EC.

HPAI has never been reported or diagnosed in poultry in Norway.

Aims

The aim of the national surveillance programme for AI in poultry is to document that the various poultry populations in Norway are free of influenza A virus of sub-types H5 and H7 and to contribute to the maintenance of this status.

Materials and methods

The programme in 2014 consisted of serological screening of blood samples from poultry. Poultry deemed at risk for exposure to influenza type A were preferentially sampled as outlined in EU's AI Directive Annex I. The sample selection was based upon a risk assessment published by the Norwegian Veterinary Institute in February 2006 (2). The sample selection included chickens, turkeys and ducks.

In addition to the samples taken from farms on the basis of the risk assessment (2), samples from breeding flocks were also tested for AI.

Blood samples were collected from at least 10 birds per holding. An exception to this was waterfowl and quail flocks, from which 50 samples were collected. If there were more than one shed on the holding, all sheds were sampled. In those instances where the flock size was less than the number required, all birds in the flock were sampled.

Chickens and turkeys

An indirect ELISA kit produced by IDEXX Laboratories, Westbrook, Maine, USA (IDEXX AI Ab Test) was used for the testing of antibodies against influenza A virus. The test has been demonstrated to detect antibody reactivity to 20 different subtypes of avian influenza including 14 hemagglutinin glycoproteins and the H5N1 subtype. The test is only validated for use in chickens.

In the case of a positive test, the sample was further examined for the presence of H5 or H7 with a haemagglutination inhibition test as described in the OIE diagnostic manual (3).

Other poultry

All serum samples from species other than chicken and turkeys were tested for specific antibodies against both H5 and H7 with the haemagglutination inhibition test.

Results and Discussion

Table 1 shows the number of flocks and birds tested in 2014.

With respect to the size of the Norwegian population of commercial poultry, an adequate number of flocks were sampled. No hobby flocks were tested in 2014 as they are not considered to be a significant population in Norway. In addition, a number of samples taken for the purposes of diagnosing disease, production problems and the control of imported animals were also screened for antibodies against Influenza A virus or H5/H7. All were negative.

Table 1. Number of certified breeder flocks, commercial flocks, hobby flocks and birds tested in the surveillance programme for AI in poultry in 2014.

Species	Certified breeder flocks		Commercial flocks		Hobby flocks		Total
	Flocks	Animals	Flocks	Animals	Flocks	Animals	Animals
Chicken	115	1 150	50	495	0	0	1 645
Turkey	5	50	45	451	0	0	501
Duck	5	250	12	550	0	0	800
Goose	0	0	2	101	0	0	101
Quail	0	0	1	50	0	0	50
Ratites	0	0	0	0	0	0	0
Total	125	1 450	110	1 647	0	0	3 097

References

1. Webster RG, Bean WJ, Gorman OT, Chambers TM, Kawaoka Y. Evolution and ecology of influenza A viruses. *Microbiol Rev* 1992; 56: 152-79.
2. Gjevre A-G, Handeland K, Jansen PA, Lyngstad TM, Ytrehus B. Risiko for smitte med høypatogen aviær influenza (HPAI) H5N1 fra ville fugler til fjørfe i Norge [Risk of transmission of HPAI H5N1 into the Norwegian poultry population via wild migratory birds, No]. Special report 2006. Oslo: National Veterinary Institute; 2006.
3. Office International des Epizooties. Manual of standards for diagnostic tests and vaccines for terrestrial animals (mammals, birds and bees). Vol 1. 5th ed. Paris: Office International des Epizooties; 2004. Updated version for AI from May 2005: http://www.oie.int/eng/normes/mmanual/A_00037.htm

The Norwegian Veterinary Institute (NVI) is a nationwide research institute in the fields of animal health, fish health, and food safety. The primary mission of the NVI is to give research-based independent advisory support to ministries and governing authorities. Preparedness, diagnostics, surveillance, reference functions, risk assessments, and advisory and educational functions are the most important areas of operation.

The Norwegian Veterinary Institute has its main laboratory in Oslo, with regional laboratories in Sandnes, Bergen, Trondheim, Harstad og Tromsø, with about 360 employees in total.

www.vetinst.no



Veterinærinstituttet
Norwegian Veterinary Institute

The Norwegian Food Safety Authority (NFSA) is a governmental body whose aim is to ensure through regulations and controls that food and drinking water are as safe and healthy as possible for consumers and to promote plant, fish and animal health and ethical farming of fish and animals. We encourage environmentally friendly production and we also regulate and control cosmetics, veterinary medicines and animal health personnel. The NFSA drafts and provides information on legislation, performs risk-based inspections, monitors food safety, plant, fish and animal health, draws up contingency plans and provides updates on developments in our field of competence.

The NFSA comprises three administrative levels, and has some 1300 employees.

The NFSA advises and reports to the Ministry of Agriculture and Food, the Ministry of Fisheries and Coastal Affairs and the Ministry of Health and Care Services.

www.mattilsynet.no

