



The surveillance programme for *Brucella abortus* in cattle in Norway in 2020



The surveillance programme for *Brucella abortus* in cattle in Norway in 2020

Authors

Thea B. Klem, Attila Tarpai, Siv Klevar

Suggested citation

Klem, B Thea, Tarpai, Attila, Klevar, Siv. The surveillance programme for *Brucella abortus* in cattle in Norway in 2020. Surveillance program report. Veterinærinstituttet 2021. © Norwegian Veterinary Institute, copy permitted with citation

Quality controlled by

Merete Hofshagen, Director of Animal Health, Animal Welfare and Food Safety, Norwegian Veterinary Institute

Published

2021 on www.vetinst.no
ISSN 1890-3290 (electronic edition)
© Norwegian Veterinary Institute 2021

Commissioned by / In collaboration with

Norwegian Food Safety Authority



Colophon

Cover design: Reine Linjer
Cover photo: Colourbox
www.vetinst.no

Content

Summary 3
Introduction 3
Aim 3
Materials and methods 4
Results and discussion..... 4
References..... 5

Summary

Blood samples from 128 cattle with abortion in the second half of the pregnancy from 43 different herds were analysed for antibodies against *Brucella abortus*. All the samples were negative for *B. abortus*.

Introduction

Eradication of bovine brucellosis in Norway was achieved in 1950 (1). Since 1994, the EFTA Surveillance Authority (ESA) has recognised Norway as a state officially free from brucellosis as described in ESA Decision 66/94/COL, later replaced by ESA Decision 227/96/COL.

A surveillance programme for *Brucella abortus* was launched in 2000. All samples were negative in 2000, 2001, 2003 and 2004. In 2002 however, two bulk milk samples were antibody positive. Further investigation did not confirm the positive results and it was concluded that the positive serological results most likely were false positive reactions. Since 2005 the programme has consisted of passive clinical surveillance. From 2004 to 2014 both aborted fetuses and blood samples from the dams were examined for *B. abortus* infection. After 2014 only blood samples from the dams are analysed for antibodies against *B. abortus*.

The Norwegian Food Safety Authority is responsible for implementing the surveillance programme for *Brucella abortus*. The Norwegian Veterinary Institute is in charge of planning the programme, performing the analyses and reporting the results. The samples are collected by inspectors of the Norwegian Food Safety Authority.

Aim

The aim of the programme is to document freedom from *B. abortus* in cattle according to demands in Directive 64/432/EEC with amendments, and to contribute to the maintenance of the present favourable situation.

Materials and methods

Herd criteria for submission of clinical material are:

- abortions occurring between the fifth month of pregnancy and 14 days before expected birth.
- at least two abortions within this pregnancy period the last twelve months.

Blood sampling is limited to one sample taken at least two weeks after the abortion. The blood sample is tested in duplicates for antibodies against *B. abortus* with an indirect ELISA ID Screen® Brucellosis Serum Indirect Multi-species test (ID.Vet, Grabels, France). If the result is doubtful or positive, the sample is re-tested in duplicates using the same ELISA. If the result then is negative, the sample is concluded to be negative. If the result is inconclusive, new blood sample from the suspected animal will be requested and tested as described above.

Doubtful or positive samples in ELISA tests are subjected to a complement fixation test (CF). If the CF test is negative the sample is concluded to be negative for antibodies against *B. abortus*. If the CF test is positive, the result is reported and new blood sample from the suspected animals is requested and tested. In addition there will be an immediate follow up with post-mortem and bacteriological investigations.

Results and discussion

A total of 128 blood samples from 43 herds (107 blood samples from 36 dairy herds and 21 blood samples from 7 beef herds) were analysed for antibodies against *B. abortus* in 2020 (Table 1).


In conclusion, there was no detection of antibodies to *B. abortus* in cattle examined in the surveillance program in 2020. Bovine brucellosis has not been detected in Norway since 1953 (1).

Table 1: Number of cattle examined for brucellosis in Norway 2000-2020

Year	Dairy cattle		Beef cattle		Total		Herds
	Foetuses (herds)	Cows (herds)	Foetuses (herds)	Cows (herds)	Foetuses (herds)	Cows (herds)	
2000	-	-	-	-	17 (14)	-	14
2001	21 (18)	-	0 (0)	-	21 (18)	-	18
2002	18 (17)	-	10 (6)	-	28 (23)	-	23
2003	30 (25)	-	4 (3)	-	34 (28)	-	28
2004	25 (21)	28 (19)	2 (2)	2 (2)	27 (23)	30 (21)	26
2005	16 (14)	48 (26)	8 (7)	8 (4)	24 (21)	56 (30)	31
2006	11 (11)	19 (13)	0 (0)	1 (1)	11 (11)	20 (14)	15
2007	11 (10)	14 (11)	1 (1)	1 (1)	12 (11)	15 (12)	12
2008	20 (17)	42 (19)	2 (1)	5 (2)	22 (18)	47 (21)	22
2009	14 (11)	19 (11)	5 (3)	7 (3)	19 (15)	26 (10)	15
2010	9 (8)	30 (15)	3 (3)	14 (4)	12 (11)	44 (19)	22
2011	7 (7)	42 (17)	2 (1)	10 (3)	9 (8)	52 (20)	22
2012	11 (10)	47 (20)	1 (1)	1 (1)	12 (11)	48 (21)	22
2013	37 (31)	130 (64)	7 (4)	22 (7)	44 (35)	152 (71)	72
2014	20 (18)	90 (37)	6 (5)	8 (7)	26 (23)	98 (44)	45
2015	-	103 (46)	-	10 (5)	-	113 (51)	51
2016	-	116 (55)	-	31 (7)	-	147 (62)	62
2017		99 (39)		28 (9)		127 (48)	48
2018		110 (48)		29 (8)		139 (56)	56
2019		98 (29)		6 (2)		104 (31)	31
2020		107 (36)		21 (7)		128 (43)	43

References

1. Sandvik O. Animal Health Standards in Norway. A historical perspective and assessment of the existing situation. Næss B (editor). Oslo: The Royal Ministry of Agriculture; 1994



*Scientifically ambitious, forward-looking
and collaborative- for one health!*



Veterinærinstituttet
Norwegian Veterinary Institute

Ås

Trondheim

Sandnes

Bergen

Harstad

Tromsø

postmottak@vetinst.no
www.vetinst.no