

The surveillance programme for bovine spongiform encephalopathy (BSE) in Norway in 2018



Veterinærinstituttet
Norwegian Veterinary Institute



The surveillance programme for bovine spongiform encephalopathy (BSE) in Norway 2018

Content

Summary	3
Introduction	3
Aim	3
Materials and methods	4
Programme outline	4
Implementation	4
Clinically suspected animals	4
Risk population and routine slaughtered animals	4
Results and discussion	5
Clinically suspected animals (passive surveillance)	5
Surveillance of slaughtered animals and fallen stock (active surveillance)	5
Conclusion	6
Acknowledgment	6
References	6

Authors
Ståle Sviland, Sylvie Lafond Benestad,
Attila Tarpai

Commissioned by
Norwegian Food Safety Authority



ISSN 1894-5678

© Norwegian Veterinary Institute 2019

Design Cover: Reine Linjer
Photo front page: Colourbox

Summary

Brain samples from 6,496 bovine animals from 2,849 different dairy herds and 1,381 beef cattle herds were submitted for examination in the surveillance and control programme for BSE in Norway 2018. All the 6,327 tested brain samples were negative for BSE.

Introduction

The BSE surveillance programme was initially based on passive surveillance (1998-2000), with active surveillance introduced in May 2000. In the period 1998-2000 the samples were investigated by histopathological examination. From 2001 onwards the samples were examined by an Enzyme-Linked ImmunoSorbent Assay (ELISA) method for detection of resistant prion protein (PrP^{Sc}).

The number of samples collected in each category in the period 1998 - 2018 is presented in Table 1. Except for the positive atypical case in 2015, BSE has never been detected in any of the examined animals.

Table 1. Number of collected samples for BSE examination by the Norwegian surveillance programme according to categories from 1998-2018.

Year	Clinically suspected animals	Fallen stock	Emergency slaughtered animals	Ante-mortem animals	Imported slaughtered animals	Healthy slaughtered animals*	Total
1998 - 2000	78	0	0	0	19	0	97
2001	14	1 352	7 073	2 612	88	2 400	13 539
2002	2	1 482	7 246	3 562	39	9 907	22 238
2003	2	1 936	7 334	4 107	39	10 727	24 145
2004	3	2 145	9 217	1 355	24	10 443	23 187
2005	1	2 318	8 462	102	10	10 486	21 379
2006	0	2 364	8 177	36	4	10 455	21 036
2007	0	2 213	7 304	48	9	10 000	19 574
2008	0	2 391	8 358	16	5	9 373	20 143
2009	1	2 435	8 320	27	3	9 451	20 237
2010	0	2 788	7 438	11	1	127	10 365
2011	1	3 078	7 241	23	1	7 878	18 222
2012	1	2 936	6 841	7	0	8 744	18 529
2013	1	3 239	7 887	9	4	9 421	20 561
2014	1	1 946	4 270	12	1	264	6 494
2015	0	1 793	5 088	48	1	0	6 930
2016	1	1 918	5 108	74	1	0	7 102
2017	2	1 640	5 227	89	1	0	6 959
2018	1	1 333	5 073	88	1	0	6 496**
Total	109	39 307	125 664	12 226	251	109 676	287 233

* Healthy slaughtered animals were excluded from the surveillance programme in 2010 and 2014-2018.

** Including 169 samples unsuitable for examination.

Aim

The aim of the surveillance programme is to document that the Norwegian cattle population is free from classical BSE.

Materials and methods

Programme outline

For 2018 the surveillance programme included examination of the following categories:

- clinically suspected cattle irrespective of age.
- all cattle older than 48 months of age, which have died or been culled, but not slaughtered for human consumption (fallen stock).
- all emergency slaughtered cattle older than 48 months.
- all cattle older than 48 months, with abnormal findings at ante-mortem examination, rejected for human consumption, or which died at the abattoir or during transport (referred to as ante-mortem animals).
- all slaughtered cattle with unknown age or origin irrespective of age.
- all slaughtered imported cattle from any country irrespective of age.

Implementation

The farmers were requested to report all cases of clinically suspected cattle irrespective of age, fallen stock older than 48 months and when delivering imported cattle to slaughter, to the Norwegian Food Safety Authority. The brain or head from clinically suspected cattle or a spoon sample from the medulla oblongata from fallen stock were submitted and analysed at the Norwegian Veterinary Institute, Oslo. Inspectors from the Norwegian Food Safety Authority collected the spoon samples of the medulla oblongata from the other categories at the abattoirs and sent them within 24 hours in a cool insulated container to the Norwegian Veterinary Institute in Oslo.

Clinically suspected animals

The usual protocol followed for the clinically suspected animals is that the whole brain is divided mid-sagittally into equal halves. One half is formalin-fixed and processed according to a standard routine protocol, embedded in paraffin, sectioned at 2 to 4 µm and stained with haematoxylin eosin (HE). Immunohistochemical staining for detection of PrP^{Sc} is performed on selected sections using a monoclonal anti-PrP antibody (SAF 84, courtesy of J. Grassi, CEA, France). From the non-fixed half, tissue from the obex area is analysed by ELISA for detection of PrP^{Sc} (TeSeE® SAP, Bio-Rad) and by Western blot (TeSeE® WESTERN BLOT, Bio-Rad) for detection of PrP^{Sc} as described by the manufacturer.

Risk population and routine slaughtered animals

Non-fixed brain tissue from the obex area was analysed by ELISA for detection of PrP^{Sc} (TeSeE® SAP, Bio-Rad) as described by the manufacturer. In cases with positive or inconclusive test results, the remaining half obex was fixed in 10% neutral buffered formalin, embedded in paraffin, sectioned at 4 µm, and stained with Hematoxylin and eosin (H&E stain) HE. Subsequently, the sections were analysed by immunohistochemical detection of PrP^{Sc} using the same protocol as for specimens from clinical suspects.

Brain samples were evaluated as unsuitable for examination when they were severely autolysed, the dorsal part of the obex area was partially missing, the obex was not present, or the medullar anatomy was not recognisable.

Results and discussion

In 2018 The Norwegian Veterinary Institute received samples from 6,496 cattle. Of these, 169 (2.6%) samples were unsuitable for examination (150 from fallen stock and 19 from emergency slaughter).

For 157 samples (2.4%) the herd of origin was not reported. However, it is important to note that in case of a positive test result from such a herd, the identity could be traced via the carcass number. The remaining 6,339 samples originated from 2,849 dairy cattle herds and 1,381 beef cattle herds. The mean number of examined animals per herd was 1.51.

Clinically suspected animals (passive surveillance)

In 2018, there were one investigated as clinical suspect due to abnormal behaviour and tremor. Improved methods for clinical examination to distinguish between real suspected BSE cases and cases with central nervous disease of other causes has probably resulted in few clinical suspected cases in later years. It is likely that animals with diseases related to the central nervous system have been examined either as fallen stock, emergency slaughtered animals or ante-mortem animals, and thus included in these categories.

Surveillance of slaughtered animals and fallen stock (active surveillance)

In 2013 The EU-commission decided to stop BSE testing in healthy slaughtered cattle in 25 EU-member states (decision 2013/76/EU). Norway stopped the testing of healthy slaughtered animals from 2014.

At the same time, the age limit for including cattle in the categories “Fallen stock”, “Emergency slaughter” and “Ante mortem” was increased to 48 months. This amendment has resulted in a reduced number of sampled animals in the categories fallen stock and emergency slaughter in 2014 and onwards compared to 2013.

The geographical distributions of the cattle population and the animals of different categories tested are presented in Table 2. There is a relatively good correlation between the collection of samples for fallen stock and emergency slaughtered animals from different regions and the distribution of the cattle population in the regions. A corresponding comparison related to the age distribution of sampled animals is shown in Table 3. The proportion of sampled animals above 48 months of age is much larger than the corresponding proportion in the population, due to the lower age limits in this programme (Table 3).

Table 2. Regional distribution of Norwegian cattle and the cattle sampled for PrPSc in 2018.

Region	Total population (%)	Fallen stock (%)	Emergency slaughter (%)	Ante mortem animals (%)	Clinically suspected animals (%)	Total (%)
Region Stor-Oslo	5.2	9.4	4.4	0.0	0.0	5.3
Region Øst	25.8	16.9	27.8	23.2	100.0	25.7
Region Sør og Vest	32.0	41.5	28.7	37.5	0.0	31.2
Region Midt	27.6	24.9	30.7	30.4	0.0	29.6
Region Nord	9.4	7.2	8.4	8.9	0.0	8.2
Total number	939 083	1 183	5 054	89	1	6 327*

There were 147 samples (2.3%) from cattle with unknown region. These samples are assumed to be distributed following the regional distribution of the cattle from known region within each target group.

* Excluding 169 samples unsuitable for examination. Including 1 imported cattle.

Table 3. Age distribution of Norwegian cattle and the cattle tested for PrPSc in 2018.

Age groups (months)	Total population (%)	Fallen stock (%)	Emergency slaughter (%)	Ante mortem animals (%)	Clinically suspected animals (%)	Total (%)
< 24	55.9	0.8	0.3	24.5	0.0	0.7 %
24-29	6.9	0.4	0.2	1.9	0.0	0.3 %
30-35	6.0	0.6	0.3	3.8	0.0	0.4 %
36-47	9.8	1.6	2.0	1.9	100.0	1.9 %
48-59	6.8	23.8	26.3	20.8	0.0	25.8 %
60-71	4.7	26.9	26.2	20.8	0.0	26.2 %
72-83	3.2	17.5	18.5	11.3	0.0	18.2 %
84-95	2.2	11.1	11.4	1.9	0.0	11.2 %
96-107	1.4	6.6	6.0	3.8	0.0	6.1 %
108-119	0.9	3.5	3.4	3.8	0.0	3.4 %
120-131	0.7	1.7	1.9	1.9	0.0	1.9 %
132-143	0.5	1.6	1.3	0.0	0.0	1.3 %
144-155	0.3	1.1	0.8	0.0	0.0	0.9 %
≥ 156	0.7	2.7	1.3	3.8	0.0	1.6 %
Total number	939 083	1 183	5 054	89	1	6 327*

There were 66 samples (1.0%) from cattle with unknown age. The age of these cattle are assumed to be distributed like the age distribution of the cattle with known age within each target group.

* Excluding 169 samples unsuitable for examination. Including 1 imported cattle.

Conclusion

In 2015, the first atypical case of BSE was detected in Norway. No case of classical BSE has ever been detected including the new 6,327 tests of 2018. It is considered highly probable with increasing certainty that the Norwegian cattle population never has been infected with classical BSE-agent due to few imports to Norway of cattle and products potentially infected with the BSE-agent, limited use of meat and bone meal in concentrates intended for ruminants, and the use of high temperature and pressure in the domestic production of meat and bone meal (1). This is supported by a quantitative risk assessment for BSE in Norway (2) and by the compiled results from the surveillance programme for BSE in the years 2001 to 2018.

Acknowledgment

The authors would like to thank all the technical staff from the Norwegian Veterinary Institute in Oslo for performing the analyses with excellence, as well as all persons involved in collecting the samples.

References

1. Mørk T, Bratberg B, Hopp P, Benestad S, Høgåsen H, Bruheim T. The surveillance and control programme for bovine spongiform encephalopathy (BSE) in Norway. In: Fredriksen B, Mørk T (editors). Surveillance and control programmes for terrestrial and aquatic animals in Norway. Annual report 2001. Oslo: National Veterinary Institute; 2002. p. 55-66.
2. Høgåsen HR, de Koeijer AA. Quantitative risk assessment for bovine spongiform encephalopathy in low- or zero-prevalence countries: the example of Norway. Risk Anal. 2007; 27:1105-17.


*Scientifically ambitious, forward-looking and cooperatively oriented
– for integrated health*

The Norwegian Veterinary Institute is a national research institute that operates in the fields of animal and fish health, food safety and feed hygiene; its primary task is to provide the authorities with independently generated knowledge.

Emergency preparedness, diagnostic services, monitoring, reference functions, consulting, and risk assessments are all important areas of activity. Our products and services include research results and reports, analyses and diagnoses, studies and advice.

The Norwegian Veterinary Institute's central laboratory and administration lie in Oslo, and we operate regional laboratories in Sandnes, Bergen, Trondheim, Harstad and Tromsø.

The Norwegian Veterinary Institute collaborates with a large number of national and international institutions.



Fish health



Animal health



Food safety



Oslo
postmottak@vetinst.no

Trondheim
vit@vetinst.no

Sandnes
vis@vetinst.no

Bergen
post.vib@vetinst.no

Harstad
vih@vetinst.no

Tromsø
vitr@vetinst.no

www.vetinst.no



Veterinærinstituttet
Norwegian Veterinary Institute