

The surveillance and control programme for Chronic Wasting Disease (CWD) in wild and captive cervids in Norway

Turid Vikøren

Ståle Sviland

Sylvie Lafond Benestad

Attila Tarpai

Torfinn Moldal



Editor Ann-Charlotte Karlsson

Technical editor Hanne Mari Jordsmyr

Scientific editors Hege Hellberg and Ståle Sviland

National Veterinary Institute



Annual Reports 2009

Surveillance and control programmes for terrestrial and aquatic animals in Norway

Title

The surveillance and control programme for Chronic Wasting Disease (CWD) in wild and captive cervids in Norway

Publisher

National Veterinary Institute
PO Box 750 Sentrum
N-0106 Oslo
Norway

Fax: + 47 23 21 60 01
Tel: + 47 23 21 60 00
E-mail: vipost@vetinst.no
Homepage: www.vetinst.no

Design: Hanne Mari Jordsmyr,
National Veterinary Institute

Front page photo: Processed from Colourbox

ISSN 1503-1454

Example of citation:

Vikøren T, Sviland S, Benestad SL, Tarpai A, Moldal T. The surveillance and control programme for Chronic Wasting Disease (CWD) in wild and captive cervids in Norway. Annual report 2009. In: Karlsson AC, Jordsmyr HM, Hellberg H, Sviland S (editors). Surveillance and control programmes for terrestrial and aquatic animals in Norway. Oslo: National Veterinary Institute; 2010.

© National Veterinary Institute

Any use of the present data should include specific reference to this report.

Chronic wasting disease (CWD) was not detected in any of the animals tested in 2009.

Introduction

CWD is a transmissible spongiform encephalopathy (TSE) of cervids (1, 2, 3). A few species of the family *Cervidae* are known to be naturally susceptible to the disease: mule deer (*Odocoileus hemionus*), white-tailed deer (*O. virginianus*), Rocky Mountain elk (*Cervus elaphus nelsoni*), and moose (*Alces alces shirasi*). CWD was first described as a clinical syndrome termed "chronic wasting disease" in captive mule deer in Colorado, USA in the late 1960s and subsequently identified as a TSE in 1978 (1). Chronic wasting disease is so far only diagnosed in free-ranging and captive cervids in North America, and is yet to be diagnosed in cervids in Europe.

Four cervid species are prevalent in natural populations in Norway: moose (*Alces alces*), red deer (*Cervus elaphus*), roe deer (*Capreolus capreolus*), and reindeer (*Rangifer tarandus*). Red deer predominate along the west coast, whereas moose and roe deer mainly inhabit other areas of the country. The wild reindeer live in dispersed populations in separate high mountain areas in southern Norway. The number officially hunted in 2009 was: 36,000 moose, 37,700 red deer, 30,000 roe deer, and 5,100 wild reindeer. Additionally, Norway has a semi-domestic reindeer population, mainly kept in the northern parts of the country, presently counting about 200,000 animals. There are approximately 75 deer farms in Norway. Most of the farms keep red deer, and only a few keep fallow deer (*Dama dama*).

Norway has large free-ranging populations of various cervids, a number of them grazing in regions where scrapie is detected, and a passive surveillance programme for CWD in Norwegian wild and captive cervids has been running from 2003. In addition, samples from slaughtered semi-domestic reindeer from several regions in the country have been tested for CWD. Norway performed an EC survey for CWD in cervids in 2006 and 2007 according to Commission decision 2007/182/EC. All samples were negative for CWD.

A small population of approximately 200 free-ranging musk ox (*Ovibus moschatus*, belonging to the *Bovidae*), inhabits the Dovre high mountain plateau in Mid-Norway. TSE has not been diagnosed in the musk ox, but the species was included in the programme from 2004.

Aim

The aim of the programme is to detect the possible occurrence of CWD in the Norwegian cervid population.

Materials and methods

Material

Tested animals included captive deer and wild cervids older than 18 months that died or were euthanized due to disease or injuries. Additionally, cervids older than 18 months necropsied at the National Veterinary Institute were examined for CWD. The number and species analysed for CWD in 2009 are given in Table 1. None musk oxen were sampled.

Methods

A rapid test (either TeSeE[®] Bio-Rad or TeSeE Sheep & Goat[®] ELISA, Bio-Rad) was used to screen brain samples for detection of the PrPCWD. All the samples were analysed at the National Veterinary Institute, which is the National Reference Laboratory for TSEs in Norway.

Results

None of the 53 samples analysed tested positive for CWD in the rapid test (Table 1).

Totally 22 of the tested animals were exclusively examined for CWD, and the majority was traffic killed roe deer (Table 1). The remaining 31 animals represent cases received at the National Veterinary Institute for routine necropsy. Two of the tested animals were captive red deer and one was a semi-domestic reindeer.

Table 1. The number of cervids tested in the Norwegian surveillance and control programme for Chronic wasting disease (CWD) 2009, distributed by reason for submission.

Species	Routine necropsy		TSE surveillance programme			Unspecified	Total
	Captive	Wild	Hunted	Traffic killed, found dead or euthanized Wild	Found dead or culled Captive		
Moose	-	9	-	2	-	-	11
Fallow deer	-	-	-	-	-	-	-
Red deer	1	7	-	-	1	-	9
Musk ox	-	-	-	-	-	-	-
Reindeer	1	1	-	-	-	-	2
Roe deer	-	12	1	18			31
Total	2	29	1	20	1		53

Discussion

No animals were positive for CWD in 2009. Mainly wild cervids were tested and among those roe deer from the eastern part of the country dominated.

Among the Norwegian cervid species, a higher risk for CWD can be assumed for red deer and moose since these species are among those known to be naturally susceptible to the disease (1- 4). Roe deer, reindeer and musk ox has so far not been found naturally infected with CWD.

References

1. Williams ES, Young S. Spongiform encephalopathies in Cervidae. *Rev sci tech Off int Epiz* 1992; 11: 551-567.
2. Williams ES. Chronic Wasting Disease. *Vet Pathol* 2005; 42: 530-49.
3. Baeten LA, Powers BE, Jewell JE, Spraker TR, Miller MW. A natural case of Chronic Wasting Disease in a free-ranging moose (*Alces alces shirasi*). *J Wildl Dis* 2007; 43: 309-314.
4. Kreeger TJ, Montgomery DL, Jewill JE, Schultz W, Williams ES. Oral transmission of chronic wasting disease in captive Shira's moose. *J Wildl Dis* 2006; 42: 640-5.

The National Veterinary Institute (NVI) is a nation-wide research institute in the fields of animal health, fish health, and food safety. The primary mission of the NVI is to give research-based independent advisory support to ministries and governing authorities. Preparedness, diagnostics, surveillance, reference functions, risk assessments, and advisory and educational functions are the most important areas of operation.

The National Veterinary Institute has its main laboratory in Oslo, with regional laboratories in Sandnes, Bergen, Trondheim, Harstad og Tromsø, with about 360 employees in total.

www.vetinst.no



Veterinærinstituttet
National Veterinary Institute

The Norwegian Food Safety Authority (NFSA) is a governmental body whose aim is to ensure through regulations and controls that food and drinking water are as safe and healthy as possible for consumers and to promote plant, fish and animal health and ethical farming of fish and animals. We encourage environmentally friendly production and we also regulate and control cosmetics, veterinary medicines and animal health personnel. The NFSA drafts and provides information on legislation, performs risk-based inspections, monitors food safety, plant, fish and animal health, draws up contingency plans and provides updates on developments in our field of competence.

The NFSA comprises three administrative levels, and has some 1300 employees.

The NFSA advises and reports to the Ministry of Agriculture and Food, the Ministry of Fisheries and Coastal Affairs and the Ministry of Health and Care Services.

www.mattilsynet.no

