

The surveillance and control programme for infectious bovine rhinotracheitis (IBR) and infectious pustular vulvovaginitis (IPV) in Norway 2015



Surveillance programmes for terrestrial and aquatic animals in Norway

Annual report 2015

Project managers at the Norwegian Veterinary Institute:

Ståle Sviland (Terrestrial animals)
Anne-Gerd Gjevre (Aquatic animals)
Mona Torp (Food safety)

Editor

Merete Hofshagen

Publisher

Norwegian Veterinary Institute
PO Box 750 Sentrum
N-0106 Oslo
Norway

Fax: + 47 23 21 60 95
Tel: + 47 23 21 60 00
E-mail: postmottak@vetinst.no
www.vetinst.no

ISSN 1894-5678

Title:

The surveillance programme for infectious bovine rhinotracheitis (IBR) and infectious pustular vulvovaginitis (IPV) in Norway 2015

Authors:

Johan Åkerstedt, Malin Jonsson, Tormod Mørk

Date: 2016-04-18

Front page photo: Hanne Mari Jordsmyr

Any use of the present data should include specific reference to this report.

Example of citation:

Åkerstedt J, Jonsson M, Mørk T. The surveillance programme for infectious bovine rhinotracheitis (IBR) and infectious pustular vulvovaginitis (IPV) in Norway 2015. *Surveillance programmes for terrestrial and aquatic animals in Norway. Annual report 2015*. Oslo: Norwegian Veterinary Institute; 2016.

The surveillance programme for infectious bovine rhinotracheitis (IBR) and infectious pustular vulvovaginitis (IPV) in Norway 2015

Johan Åkerstedt, Malin Jonsson, Tormod Mørk

All milk and blood samples tested in 2015 were negative for antibodies against bovine herpes virus (BHV-1).

Introduction

Infectious bovine rhinotracheitis and infectious pustular vulvovaginitis (IBR/IPV) is a disease caused by bovine herpesvirus 1 (BHV-1). The virus affects the upper respiratory tract of cattle causing purulent nasal discharge, hyperaemia of the muzzle and conjunctivitis. Signs of general illness are fever, depression, reduced appetite, abortions and reduced milk yield. The virus may also infect the genital tract and cause pustular vulvovaginitis and balanoposthitis. IBR/IPV is classified as a list B disease in Norway and is notifiable to the Office International des Epizooties.

Norway has not experienced clinical outbreaks of infectious pustular vulvovaginitis since the early 1960s, when two outbreaks were diagnosed in cattle. In 1993, animals in one single herd were found seropositive after primary testing of bulk milk samples. Clinical signs of IBR/IPV were not recorded and all animals on the farm were slaughtered. Attempts to isolate the virus from organ samples gave negative results. Contact herds and dairy herds in the same region were found serologically negative (1). Likewise, red deer that were shot in the neighbourhood during the hunting season the same year were found seronegative. Later, BHV-1 infection has not been demonstrated in Norway.

EFTA Surveillance Authority (ESA) has recognised Norway as free from IBR since 1994. Decisions concerning the additional guarantees relating to IBR/IPV for bovines destined for Norway are described in ESA Decision 74/94/COL. Maintenance of the ESA Decisions accepting the IBR-free status of Norway requires annual reports of the surveillance of the disease.

The Norwegian Food Safety Authority was responsible for carrying out the surveillance programme for IBR/IPV. The Norwegian Veterinary Institute was in charge of planning the programme, collecting the bulk milk samples from the dairies and performing the tests. Blood samples from beef herds were collected by inspectors from the Norwegian Food Safety Authority at slaughterhouses.

Aim

The aim of the surveillance programme for IBR/IPV was to document freedom from the infection in Norway according to the demands in ESA Decision 74/94/COL with amendments, and to contribute to the maintenance of this favourable situation.

Material and methods

The surveillance of cattle for IBR/IPV in 2015 included both dairy and beef herds. Bulk milk samples from the dairy herds were provided by the dairies, while beef cattle older than 24 months were sampled at 17 slaughterhouses, with a maximum of five animals per herd and day of sampling.

The target population of dairy herds consisted of all cattle herds delivering milk to dairies during the sampling period. In 2015, bulk milk samples from 1,176 randomly sampled dairy herds were tested. The target population of beef herds was all herds delivering cattle to slaughter in 2015. A total of 3,698 individual blood samples from 1,205 beef herds were analysed in pools. The sampled herds represented 18% of the Norwegian cattle herds. The number of herds in the surveillance programme for IBR/IPV in 2015 is given in Table 1.

All samples were tested for antibodies against bovine herpes virus 1 (BHV-1) using a commercial indirect enzyme-linked immunosorbent assay (ELISA; Boehringer Ingelheim Svanova, Uppsala, Sweden) at the Norwegian Veterinary Institute in Sandnes. In case of any positive or dubious results, a serum neutralization test would be performed.

Table 1. Numbers of dairy herds and beef herds within the frame of the Norwegian surveillance programme for IBR/IPV in 2015.

Herd category	Total no. of cattle herds*	No. of herds tested	Percent of herds tested of the total no. of herds
Dairy herds**	8 771	1 176	13
Beef herds***	4 131	1 205	29
Total	12 902	2 366	18

*Based on data from the Register of production subsidies as of 31 July 2015.

**Cattle herds delivering milk to dairies.

***Sampling performed at slaughterhouses.

Results

All bulk milk samples and blood samples tested in 2015 were negative for antibodies against BHV-1. Table 2 shows the results of the testing during the period from 1993 to 2015.

Table 2. Numbers of samples and positive results of the surveillance programme for IBR/IPV in the Norwegian cattle population during the period 1993-2015.

Year	Dairy herds	Beef herds ¹		No. of positive samples
	No. of bulk milk samples tested	No. of herds sampled	No. of individuals tested ²	
1993	26 642	0	0	1
1994	24 832	1 430	5 954	0
1995	25 131	1 532	9 354	0
1996	2 863	303	1 523	0
1997	2 654	2 214	16 741	0
1998	2 816	2 191	17 095	0
1999	2 930	2 382	18 274	0
2000	1 590	340	2 892	0
2001	2 564	434	3 453	0
2002	2 308	462	3 693	0
2003	1 845	449	3 901	0
2004	1 573	402	3 364	0
2005	1 919	484	4 766	0
2006	1 673	479	4 624	0
2007	1 575	412	4 241	0
2008	1 422	444	4 616	0
2009	1 315	435	5 048	0
2010	1 265	507	4 020	0
2011	1 226	1 278	4 758	0
2012	1 189	1 178	4 308	0
2013	1 042	1 167	4 079	0
2014	1 489	935	4 132	0
2015	1 176	1 205	3 698	0

¹Sampling performed in the herds prior to 2011.

²A small number of blood samples collected at slaughterhouses could originate from dairy herds.

Discussion

Up to 2008, a blocking ELISA (2), which had been evaluated in a retrospective analysis using a simulation model (3), was used for the surveillance programme for IBR/IPV. After participation in a proficiency testing scheme organized by the Veterinary Laboratories Agency Weybridge, New Haw, England, a commercial indirect ELISA replaced the previously used blocking ELISA. The new method was found better suited for testing bulk milk specifically.

In addition to the surveillance programme, all breeding bull candidates are serologically tested before entering the breeding centres, and all breeding bulls are subject to a compulsory test each year.

The results of the programme since 1993 strongly indicate that the Norwegian cattle population is free from IBR/IPV-infection (4).

References

1. Tharaldsen J, Krogsrud J, Ødegaard Ø. Påvist besetningsinfeksjon med bovint herpesvirus 1 (BHV-1) [Herd infection with bovine herpes virus (BHV-1) detected, No]. *Nor Vet Tidsskr* 1993; 105: 363-4.
2. Nylin B, Strøger U, Rønsholt L. A retrospective evaluation of a bovine herpes virus-1 (BHV-1) antibody ELISA on bulk-tank milk samples for classification of the BHV-1 status of Danish dairy herds. *Prev Vet Med* 2000; 47: 91-105.
3. Paisley LG, Tharaldsen J, Jarp J. A retrospective analysis of the infectious rhinotracheitis (bovine herpes virus-1) surveillance program using Monte Carlo simulation models. *Prev Vet Med* 2001; 50: 109-25.
4. Åkerstedt J, Jonsson M, Mørk T. The surveillance programme for infectious bovine rhinotracheitis (IBR) and infectious pustular vulvovaginitis (IPV) in Norway 2014. Surveillance programmes for terrestrial and aquatic animals in Norway. Annual report 2014. Oslo: Norwegian Veterinary Institute 2015.

The Norwegian Veterinary Institute (NVI) is a nationwide biomedical research institute and Norway's leading centre of expertise regarding biosafety in aquatic and terrestrial animals. The aim of the Institute is to become Norway's contingency centre of preparedness for One Health.

The primary mission of the NVI is to give research-based independent advisory support to ministries and governing authorities. Preparedness, diagnostics, surveillance, reference functions, risk assessments, and advisory and educational functions are the most important areas of operation. The Institute has its main laboratory in Oslo, with regional laboratories in Sandnes, Bergen, Trondheim, Harstad and Tromsø, with about 330 employees in total.

www.vetinst.no



Veterinærinstituttet
Norwegian Veterinary Institute

The Norwegian Food Safety Authority (NFSA) is a governmental body whose aim is to ensure through regulations and controls that food and drinking water are as safe and healthy as possible for consumers and to promote plant, fish and animal health and ethical farming of fish and animals.

We encourage environmentally friendly production and we also regulate and control cosmetics, veterinary medicines and animal health personnel. The NFSA drafts and provides information on legislation, performs risk-based inspections, monitors food safety, plant, fish and animal health, draws up contingency plans and provides updates on developments in our field of competence. The NFSA comprises two administrative levels, five regions in addition to the head office, and has some 1250 employees. The NFSA advises and reports to the Ministry of Agriculture and Food, the Ministry of Trade, Industry and Fisheries and the Ministry of Health and Care Services.

www.mattilsynet.no

