

## The surveillance and control programme for Chronic Wasting Disease (CWD) in wild and captive cervids in Norway

*Turid Vikøren*

*Ståle Sviland*

*Sylvie Lafond Benestad*

*Petter Hopp*

*Torfinn Moldal*



*Scientific editors Hege Hellberg and Ståle Sviland  
Norwegian Veterinary Institute*



**Surveillance and control programmes for terrestrial and aquatic animals in Norway.  
Annual report 2010**

**Editors**

Ståle Sviland and Hege Hellberg

**Publisher**

Norwegian Veterinary Institute  
PO Box 750 Sentrum  
N-0106 Oslo  
Norway

Fax: + 47 23 21 60 01

Tel: + 47 23 21 60 00

E-mail: [postmottak@vetinst.no](mailto:postmottak@vetinst.no)

[www.vetinst.no](http://www.vetinst.no)

**Front page photo:** Processed from Colourbox

ISSN 1503-1454

**Example of citation:**

Vikøren T, Sviland S, Benestad SL, Hopp P, Moldal T. The surveillance and control programme for Chronic Wasting Disease (CWD) in wild and captive cervids in Norway. In: Sviland S, Hellberg H (editors). Surveillance and control programmes for terrestrial and aquatic animals in Norway. Annual report 2010. Oslo: Norwegian Veterinary Institute; 2011. ISSN 1503-1454.

© Norwegian Veterinary Institute

**Any use of the present data should include specific reference to this report.**

# The surveillance and control programme for Chronic Wasting Disease (CWD) in wild and captive cervids in Norway

Turid Vikøren, Ståle Sviland, Sylvie Lafond Benestad, Petter Hopp, Torfinn Moldal

*Chronic wasting disease (CWD) was not detected in any of the animals tested in 2010.*

## Introduction

CWD is a transmissible spongiform encephalopathy (TSE) of cervids (1, 2, 3). A few species of the family *Cervidae* are known to be naturally susceptible to the disease: mule deer (*Odocoileus hemionus*), white-tailed deer (*O. virginianus*), Rocky Mountain elk (*Cervus elaphus nelsoni*), and moose (*Alces alces shirasi*). Chronic wasting disease occurs in free-ranging and captive cervids in North America, and has also been diagnosed in captive deer in South Korea in connection with deer imported from Canada. The disease is yet to be diagnosed in cervids in Europe.

Four cervid species are prevalent in natural populations in Norway: moose (*Alces alces*), red deer (*Cervus elaphus*), roe deer (*Capreolus capreolus*), and reindeer (*Rangifer tarandus*). Red deer predominate along the west coast, whereas moose and roe deer mainly inhabit other areas of the country. The wild reindeer live in dispersed populations in separate high mountain areas in southern Norway. The number officially hunted in 2009 was: 36,000 moose, 37,700 red deer, 30,000 roe deer, and 5,100 wild reindeer. Additionally, Norway has a semi-domestic reindeer population, mainly kept in the northern parts of the country, presently counting about 200,000 animals. There are approximately 80 deer farms in Norway, and 85% of them keep red deer, whereas the rest keep fallow deer (*Dama dama*).

Norway has large free-ranging populations of various cervids, a number of them grazing in regions where scrapie is detected, and a passive surveillance programme for CWD in Norwegian wild and captive cervids has been running from 2003. In addition, samples from slaughtered semi-domestic reindeer from several regions in the country have been tested for CWD. Norway performed an EC survey for CWD in cervids in 2006 and 2007 according to Commission decision 2007/182/EC. All samples were negative for CWD.

A small population of approximately 220 free-ranging musk ox (*Ovibus moschatus*, belonging to the *Bovidae*), inhabits the Dovre high mountain plateau in Mid-Norway. Sheep share summer range with the musk ox population. TSE has not been diagnosed in the musk ox, but the species has been included in the programme since 2004.

## Aim

The aim of the programme is to detect the possible occurrence of CWD in the Norwegian cervid population.

## Material and methods

### Material

Captive deer, wild cervids and musk oxen older than 18 months necropsied at the National Veterinary Institute were examined for CWD. Additionally, some wild cervids and musk oxen older than 18 months that died or were euthanized due to disease or injuries were sampled in the field. The number and species analysed for CWD in 2010 are given in Table 1.

### Methods

A rapid test (either TeSeE<sup>®</sup> Bio-Rad or TeSeE Sheep & Goat<sup>®</sup> ELISA, Bio-Rad) was used to screen brain samples for detection of the PrP<sup>CWD</sup>. All the samples were analysed at the National Veterinary Institute, which is the Norwegian Reference Laboratory for animal TSEs.

## Results

None of the 41 samples analysed tested positive for CWD in the rapid test (Table 1).

Totally 13 of the tested animals were exclusively examined for CWD, and the majority was traffic killed roe deer (Table 1). The remaining 28 animals represent cases received at the National Veterinary Institute for routine necropsy and three musk oxen that were necropsied and sampled in the field. Two of the tested animals were captive red deer.

Table 1. The number of cervids and musk oxen tested in the Norwegian surveillance and control programme for Chronic wasting disease (CWD) 2010, distributed by reason for submission.

Species	Routine necropsy		TSE surveillance programme				Total
	Captive	Wild	Wild			Captive	
			Hunted	Traffic killed	Found dead or culled	Found dead or culled	
Moose	-	12	-	-	1	-	13
Fallow deer	-	-	-	-	-	-	-
Red deer	2	2	-	-	1	-	5
Musk ox	-	4	-	-	-	-	4
Reindeer	-	2	-	-	-	-	2
Roe deer	-	6	-	11	-	-	17
<b>Total</b>	<b>2</b>	<b>26</b>	<b>0</b>	<b>11</b>	<b>2</b>	<b>0</b>	<b>41</b>

## Discussion

No animals were positive for CWD in 2010. Mainly wild cervids were tested.

Among the Norwegian cervid species, a higher risk for CWD can be assumed for red deer and moose since these species are among those known to be naturally susceptible to the disease (1- 4). Roe deer, reindeer and musk ox has so far not been found naturally infected with CWD.

## References

1. Williams ES, Young S. Spongiform encephalopathies in Cervidae. Rev sci tech Off int Epiz 1992; 11: 551-567.
2. Williams ES. Chronic Wasting Disease. Vet Pathol 2005; 42: 530-49.
3. Baeten LA, Powers BE, Jewell JE, Spraker TR, Miller MW. A natural case of Chronic Wasting Disease in a free-ranging moose (*Alces alces shirasi*). J Wildl Dis 2007; 43: 309-314.
4. Kreeger TJ, Montgomery DL, Jewill JE, Schultz W, Williams ES. Oral transmission of chronic wasting disease in captive Shira's moose. J Wildl Dis 2006; 42: 640-5.



The Norwegian Veterinary Institute (NVI) is a nationwide research institute in the fields of animal health, fish health, and food safety. The primary mission of the NVI is to give research-based independent advisory support to ministries and governing authorities. Preparedness, diagnostics, surveillance, reference functions, risk assessments, and advisory and educational functions are the most important areas of operation.

The Norwegian Veterinary Institute has its main laboratory in Oslo, with regional laboratories in Sandnes, Bergen, Trondheim, Harstad og Tromsø, with about 360 employees in total.

[www.vetinst.no](http://www.vetinst.no)



**Veterinærinstituttet**  
Norwegian Veterinary Institute

The Norwegian Food Safety Authority (NFSA) is a governmental body whose aim is to ensure through regulations and controls that food and drinking water are as safe and healthy as possible for consumers and to promote plant, fish and animal health and ethical farming of fish and animals. We encourage environmentally friendly production and we also regulate and control cosmetics, veterinary medicines and animal health personnel. The NFSA drafts and provides information on legislation, performs risk-based inspections, monitors food safety, plant, fish and animal health, draws up contingency plans and provides updates on developments in our field of competence.

The NFSA comprises three administrative levels, and has some 1300 employees.

The NFSA advises and reports to the Ministry of Agriculture and Food, the Ministry of Fisheries and Coastal Affairs and the Ministry of Health and Care Services.

[www.mattilsynet.no](http://www.mattilsynet.no)

