

## The surveillance and control programme for infectious laryngotracheitis (ILT) and avian rhinotracheitis (ART) in poultry flocks in Norway 2011

*Bruce David*

*Anne Bang Nordstoga*

*Petter Hopp*



# Surveillance and control programmes for terrestrial and aquatic animals in Norway

Annual report 2011

**Project managers at the Norwegian Veterinary Institute:**  
Ståle Sviland and Hege Hellberg

**Publisher**

Norwegian Veterinary Institute  
PO Box 750 Sentrum  
N-0106 Oslo  
Norway

Fax: + 47 23 21 60 01

Tel: + 47 23 21 60 00

E-mail: [postmottak@vetinst.no](mailto:postmottak@vetinst.no)

[www.vetinst.no](http://www.vetinst.no)

ISSN 1890-9973

**Title:**

The surveillance and control programme for infectious laryngotracheitis (ILT) and avian rhinotracheitis (ART) in poultry flocks in Norway 2011

**Authors:**

Bruce David, Anne Bang Nordstoga, Petter Hopp

**Date:** 16 July 2012

**Front page photo:** Processed from Colourbox

Any use of the present data should include specific reference to this report.

**Example of citation:**

David B, Nordstoga AB, Hopp, P. The surveillance and control programme for infectious laryngotracheitis (ILT) and avian rhinotracheitis (ART) in poultry flocks in Norway 2011. *Surveillance and control programmes for terrestrial and aquatic animals in Norway*. Annual report 2011. Oslo: Norwegian Veterinary Institute 2012.

# The surveillance and control programme for infectious laryngotracheitis (ILT) and avian rhinotracheitis (ART) in poultry flocks in Norway 2011

*Bruce David, Anne Bang Nordstoga, Petter Hopp*

*Surveillance in 2011 did not detect infectious laryngotracheitis (ILT) in chicken nor avian rhinotracheitis (ART) in turkeys.*

## Introduction

The Norwegian Food Safety Authority is responsible for the implementation of the surveillance and control programmes for infectious laryngotracheitis (ILT) and avian rhinotracheitis (ART) in chicken and turkey flocks, respectively. Started in 1998, these programmes are based on serological investigations. The Norwegian Veterinary Institute in Oslo is responsible for the planning, laboratory investigations and the reporting components of the programmes.

ILT is a severe respiratory disease in chickens that is seen in commercial chickens in most parts of the world, including most European countries (1). However, ILT has not been diagnosed in commercial chicken flocks in Norway since 1971, although clinical outbreaks of ILT have occurred sporadically in Norwegian hobby flocks since 1998 (2).

ART is a highly contagious infection which affects the upper respiratory passages of poultry. The disease is caused by avian pneumovirus (APV), and has been diagnosed in most countries (1) and sporadically in our neighbouring countries. ART had never been diagnosed in Norwegian poultry until the national surveillance and control programme for ART demonstrated the presence of antibodies against APV in one broiler breeder farm in 2003 and in one layer breeder farm in 2004. The diagnosis for ART in these flocks was based on serology only, like in many other countries (1). As the use of stamping out measures was unable to control the spread of the infection, chickens were excluded from the national surveillance and control programme for ART as of May 2005.

## Aims

The aim of the national surveillance and control programme for ILT in chickens and ART in turkeys is to document that the respective commercial poultry populations in Norway are free of these infections, and to contribute to the maintenance of this status.

## Materials and methods

According to the national regulations for certification of poultry breeding farms (3), blood samples from 60 birds must be taken at least once a year from every breeding flock at the farms. These blood samples are to be tested for Newcastle disease, as Norway is a non-vaccinating country. Thirty of the 60 samples from chicken and turkey flocks are included in the national surveillance and control programmes for ILT and ART, respectively. In addition, forty randomly chosen turkey flocks are sampled at slaughter for ART. Flocks with single positive reactions for antibodies against ILT-virus are followed up by repeated sampling, and if false positive results cannot be ruled out by this procedure, serum samples with a positive reaction in the ELISA-tests are submitted to the Veterinary Laboratories Agency (VLA), Weybridge, England for testing using virus neutralisation tests.

## ILT

An indirect ELISA-test produced by Synbiotics, was used for the testing of antibodies against the ILT-virus.

## ART

All serum samples were tested for specific antibodies against APV with a blocking-ELISA produced by SVANOVA, Uppsala, Sweden.

## Results

Of 2,818 blood samples analysed in the surveillance programme for ILT, 2,813 were negative. Positive serological results were obtained from 5 flocks originating from 4 holdings. Repeat testing of these flocks gave negative results. All 1,258 blood samples analysed in the surveillance programme for ART were negative except one sample which had intermediate result.

Table 1 shows the number of farms, flocks and birds tested in the different poultry production types in the national surveillance and control programme for ILT and ART in 2011.

Table 1. Number of farms, flocks and birds tested in the surveillance and control programmes for infectious laryngotracheitis (ILT) in chickens and avian rhinotracheitis (ART) in turkeys in 2011

Production line	No. of farms tested	No. of flocks tested	Total no. of birds tested	Flocks with seropositive samples
<i>Infectious laryngotracheitis</i>				
Broiler	64	82	2370	5
Layer	13	15	448	0
<i>Avian rhinotracheitis</i>				
Turkey	41	42	1,258	0

## Discussion

Antibodies against ILT are often found in samples from hobby flocks. It is thus of major importance that commercial poultry flocks are kept strictly isolated from hobby birds and backyard poultry flocks. The non-commercial bird populations are complex, and pose a problem for the control of this contagious poultry disease due to the difficulties associated with performing systematic disease surveillance in such bird populations.

## References

1. Saif YM, Barnes HJ, Glisson JR, Fadly AM, McDougald LR, Swayne DE (editors). Diseases of poultry, 11th ed. Ames: Iowa State University Press; 2003.
2. Løvland A, Tharaldsen J, Jonassen CM, Heier BT, 2004. The surveillance and control programmes for infectious laryngotracheitis (ILT) and avian rhinotracheitis (ART) in poultry flocks in Norway. In: Mørk T, Hellberg H (editors). Surveillance and control programmes for terrestrial and aquatic animals in Norway. Annual report 2004. Oslo: National Veterinary Institute; 2005. p. 116-119.
3. Forskrift om sertifisering av fjørfevirksomheter av 18.11.94 nr. 1020. (Provision concerning the certification of poultry enterprises.) <http://www.lovdato.no/for/sf/ld/xd-19941118-1020.html>

The Norwegian Veterinary Institute (NVI) is a nationwide research institute in the fields of animal health, fish health, and food safety. The primary mission of the NVI is to give research-based independent advisory support to ministries and governing authorities. Preparedness, diagnostics, surveillance, reference functions, risk assessments, and advisory and educational functions are the most important areas of operation.

The Norwegian Veterinary Institute has its main laboratory in Oslo, with regional laboratories in Sandnes, Bergen, Trondheim, Harstad og Tromsø, with about 360 employees in total.

[www.vetinst.no](http://www.vetinst.no)



**Veterinærinstituttet**  
Norwegian Veterinary Institute

The Norwegian Food Safety Authority (NFSA) is a governmental body whose aim is to ensure through regulations and controls that food and drinking water are as safe and healthy as possible for consumers and to promote plant, fish and animal health and ethical farming of fish and animals. We encourage environmentally friendly production and we also regulate and control cosmetics, veterinary medicines and animal health personnel. The NFSA drafts and provides information on legislation, performs risk-based inspections, monitors food safety, plant, fish and animal health, draws up contingency plans and provides updates on developments in our field of competence.

The NFSA comprises three administrative levels, and has some 1300 employees.

The NFSA advises and reports to the Ministry of Agriculture and Food, the Ministry of Fisheries and Coastal Affairs and the Ministry of Health and Care Services.

[www.mattilsynet.no](http://www.mattilsynet.no)

