



## The surveillance programme for methicillin resistant *Staphylococcus aureus* in pigs in Norway 2024

REPORT 12/2025

# The surveillance programme for methicillin resistant *Staphylococcus aureus* in pigs in Norway 2024

## Authors

Anne Margrete Urdahl, Madelaine Norström, Marianne Sunde, Carl Andreas Grøntvedt

## Suggested citation

Urdahl, Anne Margrete, Norström, Madelaine, Sunde, Marianne, Grøntvedt, Carl Andreas. The surveillance programme for methicillin resistant *Staphylococcus aureus* in pigs in Norway 2024. Surveillance program report. Veterinærinstituttet 2025. © Norwegian Veterinary Institute, copy permitted with citation

## Quality controlled by

Merete Hofshagen, Director of Animal Health, Animal Welfare and Food Safety, Norwegian Veterinary Institute

## Published

2025 on [www.vetinst.no](http://www.vetinst.no)  
ISSN 1890-3290 (electronic edition)  
© Norwegian Veterinary Institute 2025

## Commissioned by

Norwegian Food Safety Authority

## Colophone

Cover photo: Colourbox  
[www.vetinst.no](http://www.vetinst.no)



# Content

Summary ..... 3

Introduction ..... 3

Aims ..... 4

Materials and methods ..... 5

Results and Discussion ..... 5

Acknowledgement ..... 6

References..... 6

## Summary

The surveillance programme in 2024 did not detect any pig herds with MRSA. In total, 626 herds were included in the survey, of which 66 were genetic nucleus or multiplier herds, 11 herds were central units of the sow pool herds, 21 were of the largest farrow to grower or farrow to finish herds, and the remaining 528 were fattening herds.

## Introduction

The Norwegian Food Safety Authority (NFSA) is owner of the surveillance programme for MRSA in swine, and also perform the sampling. The Norwegian Veterinary Institute is responsible for planning, laboratory investigations and reporting of the programme.

There are several varieties of Methicillin resistant *Staphylococcus aureus* (MRSA) some of which are associated with animals (especially pigs), and are collectively referred to as LA-MRSA (livestock associated MRSA). Within a few years, LA-MRSAs have become widespread in pig populations around the world, thereby representing a risk for dissemination to the human population.

All types of MRSA can be transmitted between humans and animals. However, not all types are well adapted for establishing in other species than their original host. MRSA is not typically associated with disease in animals or healthy humans, but it is important to prevent transmission, especially to health institutions such as hospitals and nursing homes where the bacteria can cause severe infections that are difficult to treat.

The commercial Norwegian pig population is characterized by being essentially closed due to negligible imports of live pigs (1). The population is based on a pyramidal structure with genetic nucleus herds at the apex, and with a unidirectional downward trade of live animals. The number of herds with breeding sows were about 789 (preliminary numbers) during 2024, while the number of specialized fattening pig herds in 2023 were 1481 (2).

Surveys that were conducted in 2008 (3), 2011 (4) and 2012 (5) indicated a very low prevalence of MRSA-positive pig herds in Norway. MRSA belonging to the LA-MRSA clonal complex CC398 *spa*-type t034 was detected in pig samples for the first time in 2011 (anonymous study). In 2013/14, three clusters of LA-MRSA CC398 positive pig herds were detected (6), in eastern and southwestern Norway, respectively, and measures to eradicate LA-MRSA from positive pig herds were imposed. The rationale behind this strategy was to avoid the pig population becoming a reservoir of MRSA with the potential of zoonotic transmission. This has also been followed up in the Norwegian National Strategy against Antibiotic Resistance from 2015 (7) and in 2024 (8) stating as a goal that “LA-MRSA will not be established in the Norwegian pig population”.

The LA-MRSA eradication strategy includes restrictions on trade of live animals upon suspicion, depopulation of pigs in LA-MRSA positive pig herds, thorough cleaning and disinfection of premises, negative samples from the environment and mandatory down-time before restocking with pigs from MRSA negative herds. After restocking, samples are collected from animals and the environment several times to assess the effectiveness of MRSA eradication.

A comprehensive surveillance program of MRSA in the pig population was implemented in 2014. Sampling description and overview of results are described in the annual reports (9).

In total, nine pig herds have been found positive through the MRSA surveillance programme in 2014 - 2023 (Table 1). Additionally, contact tracing from positive herds or from persons have detected a total of 75 MRSA positive herds, bringing the total number of pig herds found positive for MRSA to 84 in this time period (9). In all LA-MRSA positive herds, measures to eradicate MRSA were imposed.

Table 1. Pig herds positive for methicillin resistant *Staphylococcus aureus* 2014 – 2024. Table show total number of MRSA positive herds detected by the MRSA surveillance programme, total number of MRSA positive herds, as well as results from the MRSA typing.

Year	No. MRSA positive herds detected by the MRSA surveillance prog. (Total no of positive herds)	Herds investigated in the MRSA surveillance prog.	MRSA typing*
2014	1 (5)	986	CC398 t034 (2), CC398 t011 (3)
2015	4 (34)	821	CC398 t034 (25), CC1 t177 (9)
2016	1 (8)	872	CC398 t034 (8)
2017	2 (6)	826	CC7 t091 (2), CC8 t024 (2), CC130 t843 (1), CC425 t6292 (1)
2018	0	716	
2019	1 (9)	722	CC398 t034 (3), CC398 t011 (5), CC130 t843 (1)
2020	0	641	
2021	0	763	
2022	0	591	
2023	0	541	
2024	0	626	
<b>Total</b>	<b>9 (84**)</b>		<b>CC398 t034 (60), CC398 t011 (8), CC1 t177 (9), CC7 t091 (2), CC8 t024 (2), CC130 t843 (2), CC425 t6292 (1)</b>

\*\* *mecC*-gene detected for CC130 t843 and CC425 t6292, *mecA*-gene detected for the others

\*\*The total number of positive herds also includes 22 herds detected during 2013

## Aims

The objective of the surveillance programme is to identify methicillin resistant *Staphylococcus aureus* (MRSA) positive pig herds with the intention of contract tracing and eradication of LA-MRSA, as the overall goal is to keep the Norwegian pig population free of LA-MRSA.



## Materials and methods

In 2024, the following were to be sampled by the NFSA: all of the genetic nucleus and multiplier herds ( $n = 72$ ) and the central unit herds of the sow pools ( $n=11$ ), the 25 largest farrow to grower or farrow to finish herds with more than 10 sows according to the Registry of Production subsidies as of 1st of October 2023, and herds with more than 70 fattening pigs ( $n = 799$ ) according to the Registry of Deliveries of carcasses for July 2023 to June 2024. Each local NFSA is expected to use their local knowledge and add herds missing in the lists, as the production subsidies registry is not complete. The genetic nucleus and multiplier herds, as well as the sow pool herds and the largest commercial sow herds, were to be sampled twice. Sampling was conducted throughout the whole year.

Pigs were sampled by using sterile SodiBox™ cloths moistened with sterile saline water. A part of the cloth was rubbed firmly against the skin behind both ears of the pig (about 5x5 cm on each side). Each cloth was used for 20 pigs, and a total of three cloths, representing 60 pigs distributed on all rooms and all age groups (except suckling piglets), were used per herd. The three cloths were analyzed as one pooled sample. In addition, in each herd two cloths were used for environmental samples taken in all rooms with pigs. Each cloth was used on about 15 control points (about 10x10 cm per location) representing furnishings, feeders, water nipples, window sills, door handles, tools, boots, ventilation system etc. These two cloths were analyzed as one pooled sample.

The samples were submitted to the Norwegian Veterinary Institute and analysed for MRSA by enrichment in 300 mL Mueller Hinton broth with 6.5% NaCl at 37°C for 18-24 h. From the culture obtained in the Mueller Hinton Broth, 10 µL were streaked on Brilliance™ MRSA2 Agar (Oxoid) and incubated at 37°C for 18-24 h. The 95% confidence interval (CI) was calculated based on a binomial distribution.

## Results and Discussion

Samples were received from 35 and 31 genetic nucleus and multiplier herds, respectively. From these 66 herds, samples were received three times from two herds, twice from 49 herds and once from 15 herds.

Samples were received from 11 sow pool herds. One herd was sampled once, ten herds were sampled twice.

Samples were received from 21 herds stated by the NFSA to be of the largest farrow to grower or farrow to finish herds, of which ten were sampled twice and eleven were sampled once.

Samples were received from a total of 528 fattening herds (out of 799).

Altogether 626 herds were included in the 2024 MRSA surveillance. This constitute a total coverage of 69.0%.

MRSA was not detected in samples from any of the 626 herds included in the 2024 surveillance (95% CI [0-0.59]).

Table 2. Number of pig herds included in the MRSA surveillance per Norwegian Food Safety Authority (NFSA) Region in 2024.

NFSA Region	No. genetic nucleus and multiplier herds	No. sow pool herds	Large Farrow to finish herds	Slaughter herds
Stor-Oslo	6	2	1	65
Øst	24	4	3	205
Sør og Vest	15	4	7	133
Midt	15	2	7	85
Nord	6	-	3	40
Total	66	11	21	528

## Acknowledgement

The authors would like to thank the technical staff for performing the analyses with excellence. In particular, Hilde Welde, has contributed substantially to the project. Moreover, the authors would like to thank all personnel from the Norwegian Food Safety Authority for collection and submission of samples.

## References

1. Årsmelding 2022, KOORIMP OG KIF. Animalia 2023. [koorimp-armelding-2022-web.pdf \(animalia.no\)](#)
2. Kjøttets tilstand 2024. Animalia 2024. [kjottets-tilstand-2024\\_30okt2024.pdf \(animalia.no\)](#)
3. NORM/NORM-VET 2008. Usage of Antimicrobial Agents and Occurrence of Antimicrobial Resistance in Norway. Tromsø / Oslo 2009. ISSN: 1502-2307 (print) / 1890-9965 (electronic).
4. NORM/NORM-VET 2011. Usage of Antimicrobial Agents and Occurrence of Antimicrobial Resistance in Norway. Tromsø / Oslo 2012. ISSN:1502-2307 (print) / 1890-9965 (electronic).
5. NORM/NORM-VET 2012. Usage of Antimicrobial Agents and Occurrence of Antimicrobial Resistance in Norway. Tromsø / Oslo 2013. ISSN:1502-2307 (print) / 1890-9965 (electronic).
6. Grøntvedt CA, Elstrøm P, Stegger M, Skov RL, Andersen PS, Larssen KW, Urdahl AM, Angen Ø, Larsen J, Åmdal S, Løtvedt SM, Sunde M, Bjørnholt JV. MRSA CC398 in humans and pigs in Norway: A "One Health" perspective on introduction and transmission. Clin Infect Dis. 2016 Dec 1; 63(11):1431-1438.
7. Nasjonal strategi mot Antibiotikaresistens 2015-2020. [https://www.regjeringen.no/contentassets/5eaf66ac392143b3b2054aed90b85210/strategi\\_antibiotikaresistens\\_230615.pdf](https://www.regjeringen.no/contentassets/5eaf66ac392143b3b2054aed90b85210/strategi_antibiotikaresistens_230615.pdf)
8. Nasjonal én-helse strategi mot antimikrobiell resistens 2024-2033. [Nasjonal én-helse strategi mot antimikrobiell resistens 2024–2033 - regjeringen.no](#)
9. The surveillance programme for methicillinresistant *Staphylococcus aureus* in pigs. Annual reports available from [MRSA hos svin \(vetinst.no\)](#).

Scientifically ambitious,  
forward-looking  
and collaborative  
– for One Health



Norwegian  
Veterinary Institute

Ås ▪ Sandnes ▪ Bergen ▪ Trondheim ▪ Harstad ▪ Tromsø

[postmottak@vetinst.no](mailto:postmottak@vetinst.no)

[vetinst.no](http://vetinst.no)