




The surveillance programme for *Echinococcus multilocularis* in red foxes (*Vulpes vulpes*) in Norway 2021



REPORT 26 / 2022

The surveillance programme for *Echinococcus multilocularis* in red foxes (*Vulpes vulpes*) in Norway 2021

Authors

Inger Sofie Hamnes, Kristin Henriksen, Kristin S. Edgar, Øivind Øines.

Suggested citation

Hamnes, Inger Sofie, Henriksen, Kristin, Edgar, Kristin, Øines, Øivind. The surveillance programme for *Echinococcus multilocularis* in red foxes (*Vulpes vulpes*) in Norway 2021. Surveillance program report. Veterinærinstituttet 2022. © Norwegian Veterinary Institute, copy permitted with citation

Quality controlled by

Merete Hofshagen, Director of Animal Health, Animal Welfare and Food Safety, Norwegian Veterinary Institute

Published

2022 on www.vetinst.no
ISSN 1890-3290 (electronic edition)
© Norwegian Veterinary Institute 2022

Commissioned by

Norwegian Food Safety Authority



Colophon

Cover design: Reine Linjer
Cover photo: Colourbox
www.vetinst.no

Content

Summary	3
Introduction	3
Aim	4
Materials and methods	4
Results and Discussion	5
Acknowledgements	9
References.....	10

Summary

The prevalence of *Echinococcus multilocularis* was based on PCR analysis of faecal samples from 511 red foxes (*Vulpes vulpes*) collected during the licensed fox-hunting season in 2021 and 20 grey wolves (*Canis lupus*) killed in 2021. None of the samples tested positive for *E. multilocularis*, documenting that the prevalence in carnivore hosts (foxes and wolves) were below 1% at a confidence level of at least 95%.

Introduction

Echinococcus multilocularis (Figure 1), the parasite causing alveolar echinococcosis in humans, is endemic in several regions of the northern hemisphere, including eastern and central parts of Europe (1, 2). During the past decades, the prevalence of *E. multilocularis* in Europe has increased in the known endemic areas (3), and the geographic distribution has expanded to regions where the parasite appeared to be absent previously (4). Similarly, alveolar echinococcosis, the life-threatening zoonotic disease caused by the metacestode stage of this tapeworm, is increasing in prevalence in Europe. A recent European project ranked *E. multilocularis* first amongst the food-borne parasites based on public health concerns (5).

The adult tapeworm resides in the small intestine of wild carnivores (definitive hosts) such as red foxes, raccoon dogs and wolves. Domestic dogs and cats can also act as definitive hosts if they prey on infected small mammals, predominantly rodents that serve as intermediate hosts.



Figure 1: *Echinococcus multilocularis*, adult worm used for spiking of positive controls included in the PCR analyses. The sack-like uterus containing hundreds of eggs is clearly visible. Worms used as controls were inactivated by kept frozen for -75 C for several days, and subsequently stored in 70% ethanol. Professor Peter Deplazes, University of Zurich, kindly donated the depicted worm. Photo: Øivind Øines, Norwegian Veterinary Institute

In Scandinavia, the first discovery of *E. multilocularis* was on the high-arctic Norwegian islands of Svalbard (6) and in Denmark (7) in 1999. However, there was no evidence of its presence in mainland Fennoscandia (8) until its detection in Sweden in February 2011 (9). Despite analyses of more than 7600 faecal samples from foxes since 2002 (10), *E. multilocularis* has not been reported in mainland Norway.

Anthelmintic treatment of dogs, prior to import from endemic regions, is compulsory in Norway to prevent introduction of the parasite. According to the EU Directive 576/2013 on pet movement, the maintenance of this national regulation requires the documentation of an *E. multilocularis*-free status within the country in question.

Aim

The aim of the surveillance is to document the freedom of *E. multilocularis* in mainland Norway.

Materials and methods

In the *E. multilocularis* surveillance of 2021, faecal samples collected from red foxes (*Vulpes vulpes*) hunted during the licensed hunting season (i.e. January to mid-April and mid-July to late December 2021) were included. In addition to faeces from foxes, samples from wolves (*Canis lupus*) killed legally, or illegally during 2021, were tested for the presence of *E. multilocularis*.

Recruitment of hunters was done through the webpages of the Norwegian Veterinary Institute (<https://www.vetinst.no/nyheter/registrering-som-provetaker-av-rodrev>). In addition, hunters who had supplied samples for the surveillance program during previous years were invited to participate by email.

Sampling containers and detailed instructions for sampling were sent to the hunters who volunteered for the program. The samples were submitted to the laboratory with written information on sample locality, date of the sampling, sex (male or female) and estimated age of the animal (juvenile or adult) in pre-paid envelopes. All counties in Norway were included in the sampling regime.

Individual faecal samples (3 g per animal) were analysed using the sensitive DNA-fishing (magnetic capture) method combined with real-time PCR detection of *E. multilocularis* mtDNA. This procedure involves magnetic capture of biotin tagged DNA-hybridisation probes targeting a locus on the *E. multilocularis* mtDNA. The biotin attached to the hybridisation probe/target DNA-complex is coupled by a noncovalent protein-protein binding interaction to streptavidin molecules which are coated onto magnetic beads. This allow extraction of parasite mtDNA from inhibitors and other DNA in the sample, by using a magnet (11).

Detection of the *E. multilocularis* DNA was carried out by real-time PCR (11, 12). If a positive real-time PCR signal is detected, the presence of *E. multilocularis* mtDNA can be verified by an additional independent real-time PCR (12), and /or using a standard PCR targeting the *nad1* gene followed by Sanger-sequencing (13). All tests are performed in duplicates with each run including two positive control DNA samples (from adult worms) and negative controls (MilliQ water) included in each run.

The DNA-fishing method is capable of detecting *E. multilocularis* DNA originating from any developmental stage of the parasite, including worms, and eggs in high volume samples. The method is suitable for use during the patent phase of the infection when eggs are shed in the faeces. This period constitutes roughly two-thirds of the entire infection period. The MC-DNA/realtime PCR methods has been shown to be more sensitive than egg isolation by sieving followed by detection of parasite DNA using a multiplex PCR, used previously in the Norwegian surveillance program (11, 12). Validation of the current methods in our laboratory has demonstrated a sensitivity of 75% and 92% in 3 g samples spiked with 5 eggs or one worm, respectively (unpublished results).

Initially, a test sensitivity of 63% and a specificity of 100% were assumed (11). However, our internal validation has demonstrated an overall sensitivity of 77 %. For samples spiked with ≥ 10 eggs the sensitivity is 89 % which is close to the method of Isaksson et al. (12) (88% test sensitivity) (11, 14). The apparent prevalence and corresponding confidence interval were estimated using EpiTools (15), with a test sensitivity of 63% and a specificity of 100%.

Results and Discussion

In 2021, 531 faecal samples from wild carnivores were analysed for *E. multilocularis*: 511 samples from red foxes (Table 1, Figure 2) and 20 samples from wolves (*Canis lupus*) (Table 1, Figure 3). All samples tested negative for *E. multilocularis* giving an estimated apparent prevalence of 0% (0.0 - 0.7%, 95%CI).

Surveillance results were no different from earlier years. All faecal samples collected from wild carnivores in mainland Norway as part of the surveillance program in 2021, were negative by PCR for *E. multilocularis*.

According to requirements of Regulation (EU) No 2018/772, Annex I, the disease freedom status must have a pathogen-specific surveillance program designed to detect a prevalence of $\leq 1\%$ at minimum confidence level of 95%. The number of samples collected and analysed in Norway in 2021 was sufficient to document a current prevalence of *E. multilocularis* below 1%.

Table 1: Number and origin (county) of red foxes and wolves examined for *Echinococcus multilocularis* in Norway during the red fox licensed hunting season in 2021 (January to mid-April and mid-July to late December) and corresponding numbers for the period 2002 - 2020.

County 2021	County 2019	Number of red foxes tested		Other species tested 2021
		2021	Total 2002-2020	
Viken	Østfold	79	922	
	Akershus	33	784	5
	Buskerud	56	388	
Oslo	Oslo	20	168	
Innlandet	Hedmark	39	1089	12
	Oppland	33	506	1
Vestfold og Telemark	Vestfold	12	115	
	Telemark	20	353	
Agder	Aust-Agder	11	165	
	Vest-Agder	7	158	1
Rogaland	Rogaland	18	127	
Vestland	Hordaland	37	345	
	Sogn og Fjordane	19	295	1
Møre og Romsdal	Møre og Romsdal	27	192	
Trøndelag	Trøndelag	39	1010	
Nordland	Nordland	33	324	
Troms og Finnmark	Troms	24	524	
	Finnmark	4	172	
Total	Total	511	7637	20 wolves

However, it is worrying that the rising prevalence in countries close to Norway has increased the risk of introduction of the parasite to Norway. In Sweden, there are already detections of *E. multilocularis* in four different regions (10), and surveillance in Denmark has demonstrated its presence in two regions (16). Studies in Sweden have discovered *E. multilocularis* in the intermediate hosts of field vole (*Microtus agrestis*) and water voles (*Arvicola amphibious*) in two study areas (20). Moreover, studies in the Baltics have shown a wider distribution of the tapeworm than previously anticipated, which has caused an increasing number of alveolar echinococcosis cases in humans (17). This is worrying, as a lack of compliance with the anthelmintic treatment requirements for pets entering the Norway after having visited endemic areas has been demonstrated (18, 19). The above-mentioned points illustrate why it is imperative to continue with the surveillance for *E. multilocularis* in Norway to document and ensure Norway has a continuous disease-free status via the annual surveillance program.

Our results support the continuing national regulation for compulsory anthelmintic treatment of imported dogs to minimize the risk of an introduction of *E. multilocularis* to Norway.

Echinococcus multilocularis surveillance 2021 (Foxes sampled at the municipal level)




Figure 2: Map of Norway showing the origin of red foxes by municipality, tested for *Echinococcus multilocularis* during the red fox licensed hunting season for red fox in 2021.




Figure 3: Map of Norway showing the origin of wolves by municipality, tested for *Echinococcus multilocularis* in 2021.



Figure 4: Map showing observations of red fox in Norway

This is an online service where citizens can logon and register their observations of fauna and flora in Norway.

Source: Norwegian Biodiversity Information Centre <https://artsdatabanken.no/Pages/180936/Roedrev>

Eide NE (2015). Rødrev *Vulpes vulpes* (Linnaeus, 1758). www.artsdatabanken.no/Pages/180936. Downloaded 26/06/2022


Acknowledgements

The authors would like to thank the technical staff for performing the analyses with excellence. In particular, Agathe Vikre Danielsen, Saima Nasrin Mohammad, Aina Steinhaug Barstad and Chiek Er have contributed substantially to the project.

References

1. Schweiger A, Ammann RW, Candinas D, Clavien PA, Eckert J, Gottstein B, Halkic N, Meullhaupt B, Prinz BM, Reichen J, Tarr PE, Torgerson PR, Deplazes P. Human alveolar echinococcosis after fox population increase, Switzerland. *Emerg Infect Dis* 2007; 13: 878-882.
2. Eckert J, Deplazes P. Biological, epidemiological, and clinical aspects of echinococcosis, a zoonosis of increasing concern. *Clin Microbiol Rev* 2004; 17: 107-135.
3. Romig T, Dinkel A, Mackenstedt U. The present situation of echinococcosis in Europe. *Parasitol Int.* 2006;55(Suppl):S187-91 10.1016/j.parint.2005.11.028
4. Combes B, Comte S, Raton V, Raoul F, Boue F, Umhang G, Favier S, Dunoyer C, Woronoff N, Giraudoux P. Westward spread of *Echinococcus multilocularis* in foxes, France, 2005-2010. *Emerg Infect Dis* 2012, 18:2059-2062.
5. Bouwknegt M, Devleeschauwer B, Graham H, Robertson LJ, van der Giessen JWB, The Euro-FBP workshop participants Prioritisation of food-borne parasites in Europe, 2016. *Euro Surveill.* 2018; 23(9):pii=17-00161. <https://doi.org/10.2807/1560-7917.ES.2018.23.9.17-00161>
6. Dahlberg T, Evans R, Slettbakk T, Ottesen P, Blystad H. *Echinococcus multilocularis* påvist på Svalbard. MSIS-rapport 2000; 28: 23.
7. Kapel CMO, Saeed I. *Echinococcus multilocularis* - en ny zoonotisk parasit i Danmark. *DVT* 2000; 83: 14-16.
8. Wahlström H, Isomursu M, Hallgren G, Christensson D, Cedersmyg M, Wallensten A, Hjertqvist M, Davidson RK, Uhlhorn H, Hopp P. Combining information from surveys of several species to estimate the probability of freedom from *Echinococcus multilocularis* in Sweden, Finland and mainland Norway. *Acta Vet Scand* 2011, Feb 11;53:9.
9. Osterman LE, Juremalm M, Christensson D, Widgren S, Hallgren G, Ågren EO, Uhlhorn H, Lindberg A, Cedersmyg M, Wahlström H. First detection of *Echinococcus multilocularis* in Sweden, February to March 2011. *Euro Surveill.* 2011;16(14):pii=19836.
10. Wahlström H, Enemark HL, Davidson RK, Oksanen A. Present status, actions taken and future considerations due to the findings of *Echinococcus multilocularis* in two Scandinavian countries. *Vet Parasitol* 2015, 213:178-181.
11. Øines Ø, Isaksson M, Hagstöm Å, Tavoranpanich S and Davidson RK. Laboratory assessment of sensitive molecular tools for detection of low levels of *Echinococcus multilocularis*-eggs in fox (*Vulpes vulpes*) faeces. *Parasites & Vectors* 2014, 7:246.
12. Isaksson M, Hagström Å, Armua-Fernandez MT, Wahlström H, Ågren EO, Miller A, Holmberg A, Lukacs M, Casulli A, Deplazes P and Juremalm M. A semi-automated magnetic capture probe based DNA extraction and real-time PCR method applied in the Swedish surveillance of *Echinococcus multilocularis* in red fox (*Vulpes vulpes*) faecal samples. *Parasites & Vectors* 2014, 7:583.
13. Trachsel D, Deplazes P, Mathis A, 2007. Identification of taeniid eggs in the faeces from carnivores based on multiplex PCR using targets in mitochondrial DNA. *Parasitology* 134, 911-920.
14. Davidson RK, Øines Ø, Madslie K, Mathis A. *Echinococcus multilocularis* - adaptation of a worm egg isolation procedure coupled with a multiplex PCR assay to carry out large scale screening of red foxes (*Vulpes vulpes*) in Norway. *Parasitol Res* 2009; 104 (3): 509-514.
15. Epitools - Epidemiological calculators. Ausvet©2020. <https://epitools.ausvet.com.au/trueprevalence>
16. Petersen HH, Al-Sabi MNS, Enemark HL, Kapel CMO, Jørgensen JA, Chriel M. *Echinococcus multilocularis* in Denmark 2012-2015: high local prevalence in red foxes. *Parasitol Res* 2018 <https://doi.org/10.1007/s00436-018-5947-y>
17. Marcinkuté A, Šarkunas M, Moks E, Saarma U, Jokelainen P, Bagrade G, Laivacuma S, Strupas K, Sokolovas V, Deplazes P. *Echinococcus* infections in the Baltic region. *Vet Parasitol* 2015, 213: 121-131.

18. Davidson RK, Robertson LJ. European pet travel: misleading information from veterinarians and government agencies. *Zoonoses Public Health* 2012, 59: 575-583.
19. Hamnes IS, Klevar S, Davidson RK, Høgåsen HR, Lund A. Parasitological and serological investigation of samples from stray dogs imported into Norway from Eastern European countries [in Norwegian]. In Norwegian Veterinary Institute's Report Series: report 15 (Norwegian Veterinary Institute), p 20.
20. Miller AL, Olsson GE, Walburg MR, Sollenberg S, Skarin M, Ley C, Wahlström H, Höglund J. First identification of *Echinococcus multilocularis* in rodent intermediate hosts in Sweden. *Int J Parasitol Parasites Vildl.* 2016. 5(1): 56-63. doi: 10.1016/j.ijppaw.2016.03.001



*Scientifically ambitious, forward-looking
and collaborative- for one health!*



Veterinærinstituttet
Norwegian Veterinary Institute

Ås

Trondheim

Sandnes

Bergen

Harstad

Tromsø

postmottak@vetinst.no
www.vetinst.no